# MODERN MEDICINE

The learned of Diagnosis and Treatment



in William & Hell (er jap s)

NOW...

## B12 activity, orally,

IN A BLOOD-BUILDING,
APPETITE-BUILDING IRON TONIC!

- B<sub>12</sub> activity of at least 12 micrograms of vitamin B<sub>12</sub> per oz. as determined by microbiological assay.
- · Iron (ferrous gluconate) in hematinic quantities.
- B complex vitamins well in excess of known minimum daily requirements.
- · Pleasant tasting, too!

PLIVID

CAPSULES

## BETA-CONCEMIN FERRATED

IRON B COMPLEX WITH BIZ ACTIVITY



Reis Concenia &



In Kondremul, each micro-globule is coated with a tough film of chondrus which resists gastrointestinal enzymic action—yet Kondremul pours freely, from the bottle, is of velvety softness.

KONDREMUL, being finely subdivided, contributes soft bulk to the dry fecal residue, easing elimination and encouraging regular bowel habits.

KONDREMUL Plain (containing 55% mineral oil).

KONDREMUL with non-bitter Extract of Cascara (4.42 Gm. per 100 cc.)

**KONDREMUL** with Phenolphthalein — .13 Gm. (2.2 grs.) per tablespoonful.

Kondremul

AN EMULSION OF MINERAL OIL AND IRISH MOSS

# Also in tablet form

—the original Irish Moss—Methyl Cellulose Bulk Laxative in Tablet Form.

KONDRETABS induce soft, easily eliminated bulk—no bloating, griping, impaction. Convenient, pleasant, easy to take.

THE E. L. PATCH COMPANY

STONEHAM, MASSACHUSETTS



"The minor and major, overt and hidden, temporary and continuing effects on the human body and mind of insufficiency of iodine in the diet . . . "\*



A Demonstrably Superior Iodine Preparation For Internal Use

Recent editorial comment\* emphasizes the importance of providing enough iodine in the diet, citing endemic iodine deficiency and trade experience showing substantial loss of iodine from table salt during transportation and storage. A long-range program is recommended to make fully potent iodized table salt universally available to the exclusion of the noniodized variety.

Elevation of blood pressure, with nervous excitement, sleeplessness, tremor and tachycardia may be induced, at least in part, through marginal, often unrecognized deficiencies of iodine. In such conditions empiric administration of iodine may prove beneficial. And in frank hyperthyroidism, iodine therapy is of course definitely indicated.

Organidin (Wampole) is an exceptionally well tolerated, unique preparation of iodine organically combined by reaction with glycerin for internal administration, entirely free of inorganic iodides, negative to starch test solution, and standardized to contain 2.5 Gm. of iodine per 100 cc. Bottles of 30 cc. with dropper (1 minim per drop). Samples and literature on request.

\*Editorial Comment: N.Y. State J. Med.: 2770, 1949.



HENRY K. WAMPOLE & CO., INCORPORATED

Manufacturing Pharmacists Since 1872
PHILADELPHIA 23. PA.

## going deep

The "hyperkinemic" activity of Baume Bengué goes beneficially deep.

It enhances blood flow through the tissue area in arthritis, myositis, muscle sprains, bursitis and arthralgia. As Lange and Weiner¹ determined by the use of thermo-needles, hyperkinemic effect may extend to a depth of 2.5 cm.

Baume Bengué also promotes systemic salicylate action. It provides the high concentration of 19.7% methyl salicylate (as well as 14.4% menthol) in a specially prepared lanolin base to foster percutaneous absorption.

Baume Bengué

Lange, K., and Weiner, D.: J.
 Invest. Dermat. 12:263 (May) 1949.

Thos. Leeming & Co. Inc. 155 E. 44th St., New York 17, N.Y.

# MODERN MEDICINE

THE JOURNAL OF DIAGNOSIS AND TREATMENT

#### EDITORIAL BOARD

James T. Case, M.D., Chicago Franklin D. Dickson, M.D., Kansas City Julius Hess, M.D., Chicago Walter B. Hoover, M.D., Boston Foster Kennedy, M.D., New York City John C. Krantz, Jr., Ph.D., Baltimore A. J. Lanza, M.D., New York City Milton S. Lewis, M.D., Nashville George R. Livermore, M.D., Memphis Francis W. Lynch, M.D., St. Paul Cyril M. MacBryde, M.D., St. Louis Karl A. Meyer, M.D., Chicago Alton Ochsner, M.D., New Orleans Robert F. Patterson, M.D. Knoxville Edwin B. Plimpton, M.D., Los Angeles Fred W. Rankin, M.D., Lexington, Ky. John Alton Reed, M.D., Washington Rufus S. Reeves, M.D., Philadelphia Leo Rigler, M.D., Minneapolis Dalton K. Rose, M.D., St. Louis Howard A. Rusk, M.D., New York City Roger S. Siddall, M.D., Detroit James S. Simmons, M.D., Boston W. Calhoun Stirling, M.D., Washington Frank P. Strickler, M.D., Louisville Richard Torpin, M.D., Augusta, Ga. Robert Turell, M.D., New York City Dwight L. Wilbur, M.D., San Francisco Paul M. Wood, M.D., New York City Irving S. Wright, M.D., New York City

SCIENCE WRITERS: Gene M. Lasater, M.D., Norman Shrifter, M.D., T. C. Papermaster, M.D., Paul D. Erwin, M.D., Bernardine Lufkin, Elizabeth Kane, Shanna McGee.

### EDITORIAL CONSULTANTS

E. R. Anderson, M.D. Joe W. Baird, M.D. William C. Bernstein, M.D. Lawrence R. Boies, M.D. Edward P. Burch, M.D. James B. Carey, M.D. C. D. Creevy, M.D. C. J. Ehrenberg, M.D. W. K. Haven, M.D. Ben I. Heller, M.D. Miland E. Knapp, M.D. Ralph T. Knight, M.D. Elizabeth C. Lowry, M.D. John F. Pohl, M.D. Wallace P. Ritchie, M.D. M. B. Sinykin, M.D. A. V. Stoesser, M.D. Arthur L. H. Street, LL.B. Marvin Sukov, M.D. Harry A. Wilmer, M.D.

#### EDITORIAL STAFF

A. E. Hedback, M.D. Editor

Thomas Ziskin, M.D. Associate Editor

Maurice B. Visscher, M.D. Consultant in Sciences

Mark S. Parker Executive Editor

Sarah A. Davidson Managing Editor

Daisy Stilwell Art Editor

James Niess Editorial Board Secretary

Lorraine Hannon Mary Worthington Editorial Assistants



## Feosol Tablets are the standard iron therapy

In simple irondeficiency anemias, Feosol Tablets are standard because they are not only easily tolerated but also remarkably effective.

Feosol Tablets' special coating assures
timed disintegration in the acid medium of
the stomach and upper duodenum where
iron is best absorbed. That is one reason why they
are standard in hospitals and with leading hematologists.
Each Feosol Tablet contains 3 grains exsiccated ferrous sulfate,


Each Feosol Tablet contains 3 grains exsiccated ferrous sulfate equivalent to approximately 5 grains crystalline ferrous sulfate.

Smith, Kline & French Laboratories, Philadelphia 'Feosol' T.M. Reg. U.S. Pst. Off.

SAVE

time, motion, space

with disposable CARTRIDS



in a permanent metal syringe

Immediately usable, premeasured doses in Cartrids eliminate the bother of filling and simplify the problems of sterilization and of storage space in the physician's bag or office.

Detailed information and literature on Cartrids 'Duracillin A.S.' and Cartrids Dihydrostreptomycin Sulfate, Crystalline, Solution are personally supplied by your Lilly medical service representative or may be obtained by writing to

ELI LILLY AND COMPANY . INDIANAPOLIS 6, INDIANA, U.S.A.

\* of either 'Duracillin A.S.' (Procaine Penicillin-G in Aqueous Suspension, Lilly), 300,000 units

or Dihydrostreptomycin Sulfate, Crystalline, Solution, 0.5 Gm. in 1 cc.

Lilly

## TABLE of CONTENTS

LETTER FROM THE EDITOR	16
CORRESPONDENCE	18
QUESTIONS & ANSWERS	26
FORENSIC MEDICINE	34
WASHINGTON LETTER	48
MEDICINE	
Clinical Physiology of Potassium William S. Hoffman	65
Iron Deficiency Anemia Dana C. Mitchell, Jr., and O. B. Mayer	67
Procaine Amide	68
Funnel Chest	68
Atypical Pneumonia	68
Management of Bacterial Pneumonias  Harrison F. Flippin and  Harold L. Israel	69
Liver Tests in Congestive Heart Failure  Leonard Felder, Alvin Mund,  and Julius G. Parker	72
Hematology in Atomic Bomb Syndrome George V. LeRoy	73
Rapid Warming and Prolonged Massage for Frostbite Wayne G. Brandstadt	74
PICTURE REPORT	



for February 15 1951

Modern Medicine Vol 19. No. 4

THE MAN ON THE COVER is Dr. William S. Hoffman, Director of Biochemistry at the Hektoen Institute for Medical Research of the Cook County Hospital, Chicago. He is also Professorial Lecturer in Physiology at the University of Illinois College of Medicine. Before assuming the directorate of the Hektoen Institute, Dr. Hoffman was a member of the National Research Council, and served on the medical faculty at Johns Hopkins University and the University of Chicago medical schools. In addition to his administrative and investigative activities he is a frequent contributor to medical journals. The article "Clinical Physiology of Potassium" on page 65, adapted from a report with the same title appearing in the Journal of the American Medical Association, is one of Dr. Hoffman's most recent contributions.

Burns Unit in British Hospital Gets Results 75





Contents for February 15 1951

CONTINUED

#### SURGERY Operative Treatment of Gigantic Hernias Sidney E. Ziffren and Nathan A. Womack...... 87 Surgical Management of Indurated Leg Hemorrhage from Esophageal Varices Howard K. Gray and Frank B. Whitesell, Jr.......... 91 Odors from Colostomy.......... 92 Biliary Infection...... 92 SURGICAL TECHNIGRAM Lumbar Sympathectomy OBSTETRICS Analgesia in First Stage of Labor Archie A. Abrams ..... 99 GYNECOLOGY Vaginal Hysterectomy......100 Trichomonas Vaginitis.....100 OTOLARYNGOLOGY Poliomyelitis after Tonsillectomy Gaylord W. Anderson, Genevieve Anderson, Audrey E. Skaar, and Fenestration Operation for Otosclerosis OPHTHALMOLOGY Diagnosis of Intracranial Aneurysms Eldridge Campbell and C. W. Burklund . . . . . . . . . . . . . . . . . . 103 DERMATOLOGY Eczema from Wooden-handled Objects 1. B. Howell and D. Shelton Blair .... 106 **ORTHOPEDICS** Intracapsular Hip Fractures Robert T. McElvenny.....108 NEUROLOGY Phenylacetylurea.....112 Neuropathy in Infectious Mononucleosis Theodore C. Bernstein and Harold G. Wolff ......114



Congestion

Advantages without disadvantages

Specify

the Original

ARGYROL Package The advantages—the <u>positive</u> factors favoring ARGYROL—are enough to make it an overwhelming choice: a rare combination of decongestive, demulcent, detergent and bacteriostatic actions.

Lack of disadvantage—the outstanding disadvantage of rebound congestion, so typical of vaso-constrictors, is absent when the choice is ARGYROL.

#### The ARGYROL Technique

- The nasal meatus . . . by 20 per cent ARGYROL installations through the nasolacrimal duct.
- The nasal passages . . . with 10 per cent ARGYROL solution in drops.
- The nasal cavities . . . with 10 per cent ARGYROL by nasal tamponage.

#### Its Three-Fold Effect

- Decongests without irritation to the membrane and without ciliary injury.
- Definitely bacteriostatic, yet non-toxic to tissue.
- Stimulates secretion and cleanses, thereby enhancing Nature's own first line of defense.

the medication of choice in treating para-nasal infection.

Made only by the

A. C. BARNES COMPANY, NEW BRUNSWICK, N. J.

ARGYROL is a registered trademark, the property of A. C. Barnes Company



Contents for February 15 1951

CONTINUED

#### GERIATRICS

#### RADIOLOGY

#### PROCTOLOGY

#### PEDIATRICS

#### UROLOGY

Surgery of Acute Cholecystitis.....130

#### MEDICAL FORUM

BASIC SCIENCE BRIEFS......156

SHORT REPORTS......160

CURRENT BOOKS & PAMPHLETS......182

PATIENTS I HAVE MET......188

Modern Medicine, The Journal of Medical Progress, is published twice monthly on the first and fifteenth of each month at Minneapolis, Minn. Subscription rate: \$5.00 a year, 25c a copy. Business Manager: M. E. Herz. Address editorial correspondence to 84 South 10th Street, Minneapolis 5, Minn. Telephone: Bridgeport 1291. Advertising Representatives: New York 17: Lee Klemmer, George Doyle, Bernard A. Smiler, 50 East 42nd Street, Suite 401. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Central 6-4619. San Francisco 4: Duncan A. Scott & Co., Mills Bldg. Telephone: Garfield 1-7950. Los Angeles 5: Duncan A. Scott & Co., 2978 Wilshire Blvd. Telephone: Dunkirk 8-4151.

## ա intractable peptic ulcer

# KUTROL

UROENTERONE, PARKE-DAVIS

#### PEPTIC ULCER INHIBITANT

## when other therapy fails

#### Beneficial response in up to 70 to 80 per cent of cases -

Noteworthy results in chronic cases of Duodenal and Jejunal Ulcers having frequent recurrences and resistant to intensive conventional therapy.

Remission of ulcer often within 3 to 6 weeks –

Rapid relief of symptoms and disappearance of ulcer crater.

#### Simplified regimen -

Normal three-meals-a-day schedule soon after treatment begins.

#### Well-tolerated -

Does not inhibit gastric secretion. No toxicity or idiosyncrasy noted.

DOSAGE: Two KUTROL Kapseals® four times daily, one-half hour before mealtime and at bedtime.

PACKAGING: KUTROL Kapseals®, 75 mg., are supplied in bottles of 100.

Descriptive Literature Available On Request.

PARKE, DAVIS & COMPANY



# SYNDROX

Brand of Methamphetamine Hydrochloride



McNEIL

## **Control of Obesity in Pregnancy**

When control of excessive weight gain during pregnancy is desirable, Syndrox attacks the problem from two angles:

## SYNDROX\* HYDROCHLORIDE

- 1. suppresses the appetite;
- 2. by elevating the mood, tends to alleviate the psychic factors frequently responsible for over-eating.

Syndrox also has been used to advantage in many conditions in which its stimulating effect on the higher cerebral centers is desirable:

MILD DEPRESSIVE STATES — produces a sense of well-being and increased energy.

**DROWSINESS, NARCOLEPSY** — counteracts abnormal sleepiness . . . can be used to overcome the sedative action of the antihistaminics.

ACUTE AND CHRONIC ALCOHOLISM—adjunctive treatment.

Syndrox acts quickly (onset 10 to 20 minutes); effect is prolonged (6 to 12 hours, depending on dose); has negligible side-effects, with proper dosage; relatively small dosage.

#### SUGGESTED DOSAGE:

Initially 2.5 to 5 mg. daily; dosage may be increased to 2.5 to 5 mg. two to three times daily and maintained at this level as long as there are no untoward effects. *Caution*: Use only as suggested.

(scored, green) in bottles of 100, 1000; also available—a pleasanttasting elixir (colored amber) each 30 cc. (1 fl. oz.) containing

Supplied in 5 mg. tablets

\*Trade-Mark of McNeil Laboratories, Inc.

Samples on request

20 mg. Pints and gallons.

#### LETTER FROM THE EDITOR

#### Dear Reader:

Since the time of Hippocrates, the doctor's chief concern has been the treatment and the care of the sick. But while the problem has remained the same, the means have changed.

The story of the progress of medical science has been told so often that we have a tendency to accept the historic fact and to take today's advances for granted. Only a dozen years ago pneumonia was one of the major causes of death. Now a patient rarely dies of pneumonia. The sulfonamides and antibiotics have given the medical profession the means of effective therapy.

Less spectacular, perhaps, but tremendously important in the treatment of other diseases are the anticonvulsant, the anticoagulant, and the antihistaminic drugs. In the past few years great strides have been made in the development and employment of these therapeutic agents to relieve suffering and to save lives.

Clinical progress with these drugs is the theme of three special articles now in preparation for *Modern Medicine*. These articles will outline the most successful modes of treatment with the three groups of drugs. Indications, efficacy, and limitations will be evaluated in the light of present knowledge.

The first article will appear in the March 1 issue. The paper, entitled "Control of Seizures with Drugs," will be a comprehensive review of the antisconvulsants by Dr. William G. Lennox, Assistant Professor of Neurology, Harvard Medical School, and President of the International League of Epilepsy since 1935.

The second article, "Anticoagulant Therapy for Thromboembolic Disease," will be published in April. The author is Dr. Irving S. Wright, Associate Professor of Medicine, Cornell University, and Chief Consultant in cardiovascular diseases to the Veterans Administration.

The third article, "The Antihistaminics" by Dr. Morris Fishbein, former Editor of the Journal of the American Medical Association, will appear in the June 1 issue.

A Nebback

announcing a superior, highly palatable sedative-antispasmodic

## elixir 'ESKAPHEN B with BELLADONNA'

Elixir 'Eskaphen B with Belladonna' combines, in a light and delightfully flavored elixir:

- 1. **All** the natural alkaloids of the time-proved antispasmodic: **belladonna**...to combat spasm.
- 2. The mild, calming sedative: phenobarbital...to relieve nervous tension and reduce reflex excitability.
- 3. Full therapeutic dosage of the virtually specific nutrient and restorative: **thiamine** . . . to help rectify dietary deficiencies.

You will find broad therapeutic application for elixir 'Eskaphen B with Belladonna' in the many spastic conditions of smooth muscle. It will be of particular value, however, in the treatment of spastic conditions of gastro-intestinal musculature.

Formula: Each 5 cc. teaspoonful contains: natural belladonna alkaloids, 0.2 mg.; phenobarbital, ¼ gr. (16 mg.); thiamine, 5 mg. (nearly three times the minimum daily requirement); alcohol, 15%. Available in 6 fl. oz. bottles.

Smith, Kline & French Laboratories, Philadelphia

'Eskaphen B' T.M. Reg. U.S. Pat. Off.

## Correspondence

Communications from the readers of Modern Medicine are always welcome. Address communications to The Editors of Modern Medicine, 84 South 10th St., Minneapolis 3, Minn.

#### Relief of Herpetic Pain

10 THE EDITORS: In the December 15, 1950 issue of Modern Medicine there is a question from a Colorado M.D. about pain following herpes zoster (p. 22). The consultant replies that no therapy is specific.

Dr. E. V. Allen and I have treated 6 patients with Etamon Chloride with 80 to 100% relief in all cases. This is by far the simplest procedure and the most effective. The usual precautions should be considered when using Etamon.

HUGH S. BROWN, M.D.

Spokane

#### Red Feather Support

TO THE EDITORS: May I extend my personal thanks for the fine support given by *Modern Medicine* to the Red Feather campaigns this fall?

You will be interested to know that final reports indicate that the Community Chests and other united Red Feather campaigns have raised record-breaking total funds. Bringing the facts to the attention of your readers undoubtedly helped a great deal in stimulating the wonderful support.

C. E. WILSON National Campaign Chairman New York City

#### Antibiotics and Biopsies

TO THE EDITORS: In a footnote to the comments on liver damage in pneumonia by Dr. Alfred R. Ross, you request your readers to write as to whether they have evidence of liver damage from antibiotic treatment of pneumonia (Nov. 1, 1950, p. 18). While I have no positive evidence on this specific point, I seriously doubt that the antibiotics have a deleterious effect upon the liver.

Dr. Ross states in his letter that liver function was not impaired in pneumonia before the advent of antibiotic treatment and that, in his opinion, such treatment causes the liver damage. I seriously question this statement of Dr. Ross and would appreciate information as to its basis. I happen to have done considerable research in the past on liver damage and repair and, from my experience, any severe infection is quite capable of producing, and frequently does produce, some degree of liver damage. I know of no reason why pneumonia should be an exception.

The article which prompted Dr. Ross's letter, "Liver in Pneumococcal Pneumonia" by Drs. H. J. Zimmerman and Lawrence J. Thomas, appeared in the September 1 number

## more rapid, complete and prolonged shrinkage with BENZEDREX INHALER:



After 5 minutes.
Benzedrex Inhaler has produced virtually complete shrinkage.



After 1 hour.
Benzedrex Inhaler still provides almost complete shrinkage.

### for comparison-shrinkage with ephedrine:



After 5 minutes.
Shrinkage does not compare with that of Benzedrex Inhaler.



After 1 hour.

The ephedrine nostril is almost completely occluded.

Before treatment the patient's inferior turbinate almost completely obstructed the airway in both cases. The only variable was the vasoconstrictor used.

These photographs reveal why physicians tell us that Benzedrex Inhaler is the best inhaler they and their patients have ever used.

You can recommend Benzedrex Inhaler for nasal congestion between office treatments with assurance. It will not cause excitation or wakefulness.

Smith, Kline & French Laboratories, Philadelphia

BENZEDREX INHALER

'Renzedrex' T.M. Rev. U.S. Pat. Off.

### Interest in life and living

When the patient settles down to "the completion of life", depression can so easily get the upper hand. The seemingly endless, daily routine of living is approached with apathy, inertia and lack of interest; and the patient's own outlook on life drags her down the path to eventual break-up—physical as well as mental.

For such a patient 'Dexedrine' Sulfate is of unequalled value. Its uniquely "smooth" antidepressant effect restores mental alertness and optimism, induces a feeling of energy and well-being—and thus has the happy effect of once again reviving the patient's interest in life and living.

Smith, Kline & French Laboratories, Philadelphia

## Dexedrine\*Sulfate

tablets • elixir •T. M. Reg. U.S. Pat. Off.

the antidepressant of choice



of Modern Medicine (p. 48). The last sentence in that article, in my opinion, is the real key to the question. The paragraph referred to is herewith quoted:

The extent of hepatic dysfunction probably depends on a combination of factors, including previous state of the liver, especially with regard to alcoholism and dietary deficiency, as well as intensity of infection, fever, anoxia, and the accompanying alarm reaction.

I, too, would be very much interested in hearing from others as to the question of antibiotics having a deleterious effect upon the liver. I personally believe that antibiotics have been blamed for a great deal of mischief for which they are not responsible.

The second item upon which I would like to comment is the article, "Indications for Biopsy of the Skin," by Dr. Earl B. Ritchie, M.D., of the University of Texas, published in Modern Medicine, November 15, 1950, p. 108. I quite agree with practically everything that Dr. Ritchie says, but he has omitted one principle which I, as a pathologist, feel is fundamental.

It has always been my belief that, unless complete surgical excision of a skin lesion is impracticable, a biopsy in the usual sense of the word should not be done. Instead of removing a portion of any questionable skin lesion for pathologic examination, the entire lesion should be widely excised so as to insure complete removal of the lesion. It has been my displeasure to receive for examination too many incompletely removed malignant lesions of the skin when complete removal would have been just as easy.

I have had this happen in the case of melanomas and other types of malignant skin lesions. Only last year a surgeon removed what clinically and histologically was a benign pigmented nevus of the face which, on the surface, was only about 1.5 mm. in diameter. The total width of the specimen was only 3 mm. Sections made through the center of this lesion showed that a considerable amount of nevus extended far beyond the limits of the section removed.

In view of the reputed tendency of benign pigmented nevi to become malignant when incompletely removed, the lesion mentioned would have been better left alone.

I feel that the principle of wide excision of suspected lesions cannot be too strongly emphasized. I have seen too many sad results from failure to so excise.

JOHN H. SCHAEFER, M.D. Los Angeles

#### Acne Rosacea and Athlete's Foot

TO THE EDITORS: I would like to submit the following discussion of two questions that appeared recently in the Questions & Answers department of Modern Medicine (Dec. 1, 1950).

The first question was concerned with the treatment of acne rosacea. Even though the etiology of rosacea is not proved beyond doubt, evidence is increasing to show that it is probably due to vitamin B deficiency. Therapeutic success seems also to support this contention. Ichthyol by oral medication (Rx ichthyoli 0.25 gm., in capsules, 1 t.i.d.) seems to



write for free Sample of Armstrong's Circle A Nurser and a supply of literature. Address Armstrong Cork Company, Drug Sundries Department, 8202 Prince Street, Lancaster, Pennsylvania.

Armstrong's A Nurser

restore the normal intestinal bacterial flora responsible for the production of the vitamin B complex. One can provide the necessary amount of vitamin B complex by giving brewers' yeast tablets (l t.i.d.). This is replacement therapy.


I do not treat locally with either type of medication. The results are excellent. My preference in most cases is Ichthyol. Roentgen therapy is dangerous; its effect is only temporary, if any. The use of hormones

has no rationale at all.

To the question, "What is the proper treatment for athlete's foot?" I would answer: Athlete's foot is not a dermatomycosis, that is, it is not caused by a fungus of any description. Fungi found in the roof of the deeply embedded, sago-grain-like blisters and pustules—the one and only primary lesion of this condition—are simple saprophytes, nosoparasites. Even they are absent in the great majority of cases.

Thus, therapeutic suggestions directed against these innocent, harmless, nosoparasitic fungi are doomed to failure, because they want to treat or attack something (fungi) which is not there at all and, even when the fungi can be found, they have nothing to do with either the etiology or pathogenesis of this condition. Without due consideration of the etiology and pathogenesis of the condition, therapeutic suggestions remain poor guesswork. However, exact knowledge of the etiology and pathogenesis has been available for a considerable length of time.

The correct scientific term of the condition under discussion is pompholyx, occurring on palms, soles, fingers, or toes, characterized by deep-seated, sago-grain-like blisters with or without pustules, embedded



\*Rehfuss, M. E.: Penna. Med. J. 42:1335, 1939.

the finger of suspicion points to

## biliary disorders

... when the patient complains of flatulence, indigestion, constipation. Every other patient past age 40 suffers from some form of biliary disturbance,\* investigators state. Caroid and Bile Salts Tablets offer simple, effective relief of dyspepsia, constipation and other distressing symptoms of biliary disorders. Functional restoration is aided by-

stimulation of bile flow

improved digestion and absorption of foods

gentle laxation without whipping the bowel

Dosage: 1 or 2 tablets after breakfast and at bedtime with a glass of water. Sample available on request

American Ferment Company, Inc. 1450 Broadway, New York 18, N. Y.

CAROID AND / BILE SALTS tablets

Specifically indicated in

biliary dyspepsia and constipation



strument ON, the Viso "does the whole job itself"—while the operator merely "steers" the recording by one control from lead to lead (Fig. 1).

No routine adjustments for sensitivity, stylus temperature, or baseline centering—once set, they stay set—only to be checked once in a while! And, never any interchanging of electrode connections. (Patients welcome this too!)

Yes, by its own design, the Viso "does the whole job" of turning out finished, permanent 'cardiograms. You simply guide its recording.

Mail coupon below for folder describing this, and many other Viso advantages.



patient connected and in-

MAIL TO	SANI	BORN	CO.	CAMBRIDGE 39, MASS
Without "Sample Cardiette	obligation Demonst	, please seration of	end me a San	new folder, bern Visc-
DR				
STREET				
CITY &	STATE.			M-2-51

in normal skin, without any inflammatory reaction, and noted by its sudden, explosion-like appearance. Pompholyx (id-eruption, dermatophytosis, or, in vulgar terms, athlete's foot) is an endoparasitic-hematogenous eruption with definite localization on hands and feet caused alone by the Bacillus endoparasiticus Benedik, 1927, as proved by mycologic, bacteriologic, and histopathologic methods. It is not infectious and not transmissible because of the existing infection immunity.

There is only one specific therapy, vaccine therapy, by the vaccine of B. endoparasiticus. No local treatment is indicated or necessary. For more details consult T. Benedek; Urol. & Cutan. Rev. 50:467-493, 1946; Indust. Med. 16:344-349, 1947; and Urol & Cutan. Rev. 54:409-413, 1950. B. Schuster, Northwest Med. 46:298, 1947.

TIBOR BENEDEK, M.D.

Chicago

#### Symposium Interesting


TO THE EDITORS: It gives me great pleasure to introduce myself to you. I am an Egyptian dermatologist, fellow of the World Health Organization, at present studying at the Dermatology Department of the University of Pennsylvania under Dr. Donald Pillsbury.

I have recently come across your special November 15 number about dermatology. I found it most interesting. I wish to have a copy of that issue.

GEORGE RIZK MIKHAIL, M.D. Philadelphia

The BEST-INFORMED Doctors read Modern Medicine regularly

in acute laryngotracheal
bronchitis
prompt response
with
Terramycin



		12 11 11 11 11 11	2000	Secretary of the
CASE				

Case report taken from Herrell, W. E.; Heilman, F. R., and Wellman, W. E.: Ann. New York Acad. Sc. 53:448 (Sept. 15) 1950.

# Terramycin HYDROCHLORIDE

Rapidly Absorbed

"... produces significant continuous blood levels when 250 mg, are given every six hours." <sup>1</sup>

Well Tolerated

"...side-effects have not constituted a problem in therapy."2

Dosage: On the basis of findings obtained in over 150 leading medical research centers, 2 Gm. daily by mouth in divided doses q. 6 h, is suggested for most acute infections.

Supplied: 250 mg. capsules, bottles of 16 and 100; 100 mg. capsules, bottles of 25 and 100; 50 mg. capsules, bottles of 25 and 100.



Antibiotic Division
CHAS. PFIZER & CO., INC.
Brooklyn 6, N.Y.

I. Weich, H.; Hendricks, F. D.; Price, C. W., and Randall, W. A.; J. A. Ph. A. (Sc. Ed.) 39:185 (Apr.) 1950. 2. Knight, V.; New York State J. Med. 50:2173 (Sept. 15) 1950.

## Questions & Answers

All questions received will be answered by letter directed to the petitioner; questions chosen for publication will appear with the physician's name deleted. Address all inquiries to the Editorial Department, MODERN MEDICINE, 84 South Tenth Street, Minneapolis 3, Minnesota.

QUESTION: Is it likely that a patient with chronic nonspecific ulcerative colitis who has had anal incontinence for the past year will eventually regain control? Are drugs helpful in restoring anal continence?

M.D., Pennsylvania

ANSWER: By Consultant in Proctology. The information given is insufficient for a definite opinion. However, a patient with ulcerative colitis with anal incontinence of a year's duration has practically no chance of regaining control. No drugs are known that would help to restore anal sphincter control. If, however, the case is one of pseudoincontinence, that is, an excessive number of loose stools with urgency or tenesmus so that the patient is unaware of bowel movements but has a sound sphincteric musculature, true continence may be expected with improvement of the intestinal disease.

QUESTION: When both parents are afflicted with severe hay fever would transference of the allergy to offspring be prevented if conception or delivery were avoided during the allergy season?

M.D., New York


ANSWER: By Consultants in Allergy and Obstetrics. Since hereditary traits are fixed within the chromo-

somes of the germ cells, the effect of external environmental factors such as active allergy should be nil. There is no evidence to indicate that planning a family by allergic parents so that conception and delivery were avoided during the hay fever season would in any way reduce allergic manifestations in the offspring. No matter when conception or birth occurs, 90% of the children of 2 allergic patients will show some form of allergy before the age of 10 years. However, there is reason for planning a family so that delivery of a child does not occur during the hay fever season, if the mother has severe hay fever, because childbirth might aggravate the condition.

QUESTION: A complete colectomy with a permanent ileostomy for chronic ulcerative colitis was done for a female patient, age 26, about a year ago. She is bothered by loud gurgling noises and an offensive odor. Have you any suggestions to mitigate these obnoxious effects?

M.D., New Jersey

ANSWER: By Consultant in Proctology. The gurgling noises are usually due to hyperperistalsis and may be reduced or eliminated by avoidance of irritating foods, such as



When dissolved slowly in the buccal sulcus...

#### LOZILLES

Tyrothricin-Propesin Lozenges

provide

## safe

... non-sensitizing

### effective

...2 mg: tyrothricin per lozenge

## topical

... no systemic absorption

## analgesic

... sustained relief

## -antibiotic

...high salivary concentrations of tyrothricin

therapy

White Laboratories, Inc., Pharmaceutical Manufacturers, Newark 7, N.J.

PRULOSE COMPLEX

### The dietary approach for Therapeutic correction of Functional Constipation.

Combining the well-known bulk-producing effect of methylcellulose with the universally accepted laxative properties of prunes, the natural laxative food, fortified with an isatin derivative. This activated moist bulk

- 1. activates the colon to normal motility
- 2. prevents drying out and hardness of the colon contents
- 3. supplies the necessary bulk to increase the volume of the stool

### These actions of PRULOSE COMPLEX

- 1. promptly relieve the symptoms of functional constipation
- 2. gently stimulate peristaltic activity
- 3. institute a return to normal colon function

#### PRULOSE COMPLEX tablets are:

- 1. convenient, small and easily swallowed
- 2. economical-low dosage
- 3. formulated for maximum patient cooperation

Send for samples and Guidance booklets for your patient

Name

Address

State

17.6

Zone

The HARROWER

Laboratory, Inc.

ACTIVATED MOIST BULK

ADULT DOSAGE: 3 or mer shiets with a full all shield daily, until rermal elimination is established; then reduce to 3 falshed; then reduce to 3 falshed; then reduce to 3 fal-

h Tublet contains: hydrated Prune Concen

athylcellulose.....16 gr.

to 19thydroxyphenyllegile 10/65 gr.) ....... (0.001 gm.

Available in bottles of 100 tablets

spices, the cabbage family, and carbonated waters. Sedatives and belladonna or belladonna derivatives, such as atropine, are helpful.

Charcoal tablets or the recently introduced Kaolin and chlorophyl tablets are effective in eliminating offensive odors. Dosage is a matter of individual adjustment. Also important is the use of a well-fitting, disposable plastic ileostomy bag. The nondisposable rubber bag retains offensive odor even after diligent washings and should be discarded.

Reassurance that the borborygmi are usually not heard by persons other than the patient helps to remove anxiety which, in itself, may accentuate the condition. Since the gurgling noises usually occur immediately.

ately after meals, the patient should be advised to take a short rest after partaking of food.

QUESTION: What time interval should intervene between the last appearance of symptoms in an epileptic patient under treatment with Dilantin and his resumption of automobile driving and work with heavy industrial machinery?

M.D., Delaware

ANSWER: By Consultant in Neurology. Most physicians treating epileptic patients believe that patients who have been free from seizures for a period of two years with medication may resume driving or work with heavy machinery with relative safety.



Royal takes extra time
to bring you Steel furniture
of enduring quality

This man is an artist with a piece of steel and a buffing wheel. His job is one of the many individualized EXTRA touches that go into the making of distinctive Royal furniture.


Today, with much of our production devoted to defense needs, and civilian demand as keen as ever . . . we could skip some of our time-taking extra operations. But we never will. Time hardly touches long-wearing Royal Furniture because Royal TAKES the time to bring you the very best.

If you cannot wait for Royal's extended deliveries, may we suggest you see our worthy competitors. Many of them make products of good quality which we recommend.



METAL FURNITURE SINCE '97

ROYAL METAL MANUFACTURING CO. 175H North Michigan Avenue • Chizago 1 New York • Los Angeles • Michigan City, Indiana Warren, Pennylvania • Preston and Galt, Ontario



for the pain that wasn't there following Pabalate therapy in arthritis.

Para-aminobenzoic acid 0.3 Gm. (5 gr.), plus sodium salicylate 0.3 Gm. (5 gr.) provide higher salicylate blood levels on lower salicylate dosage — with more prolonged clinical relief, and reduced side-effects.





Crookes now gives you a choice of two greaseless, agreeably scented, pleasant-to-apply antipru-

ENZO-CAL, A. H., new

ritic creams.

contains the outstanding antihistamine, thenylpyramine hydrochloride. On prescription only — 1 oz. tubes and 1 lb. jars.

ENZO-CAL remains unchanged, the original anesthetic (benzocaine) calamine cream so widely relied upon by the profession  $-1 \frac{1}{2}$  oz. tubes and 1 lb. jars.

Both contain soothing, protective healing colloidal calamine and zinc oxide. Both are remarkably effective in relieving itch and irritation due to ECZEMA, PRURITUS ANI ET VULVAE, EXANTHEMS, FOOD, DRUG AND PLANT RASHES, DIAPER RASH.

For SAMPLES of ENZO-CAL and ENZO-CAL, A. H.

-- write

Crookes

Crookes LABORATORIES, Inc.

305 East 45th St. N Y 17. N.Y.

QUESTION: All his life a patient, now 52, has had allergic symptoms. He reacts especially to heat. If he enters a warm room, eats warm food, or drinks warm water, tea, coffee, or milk, he immediately becomes overheated, starts to perspire, feels dizzy, and sometimes even breaks out in hives. Is there any treatment for physical allergy?

M.D., New York

ANSWER: By Consultant in Allergy. The treatment of physical allergy has never been satisfactory. Fortunately, the condition sometimes disappears spontaneously. Before the advent of the antihistaminic drugs, many clinicians recommended gradual exposure to heat or cold in the hope that tolerance could be developed. However, complete cooperation from the patient was usually difficult to obtain. Now one may try a long-acting antihistaminic drug such as Di-paralene or Perazil or a repeat action drug such as Chlor-Trimeton or laminated Pyrrolazote. Medication can be taken every eight hours.

QUESTION: A healthy woman 20 years old is concerned about the non-development of her breasts. Is there any external application to be used?

M.D., Maryland

ANSWER: By Consultant in Gynecology. If this patient has normal ovarian function as demonstrated by regular menstruation and normal pelvic findings, the use of prosthesis, commonly known as falsies, is recommended. If she has concomitant amenorrhea, the use of estrogens in large dosages given in cyclic manner to allow withdrawal bleeding would also serve to stimulate breast growth during the time of administration.



## BACITRACIN TROCHE

#### WITH BENZOCAINE

#### FOR CONTROL OF LOCAL DISCOMFORT

The clinical efficacy of bacitracin troches has been further expanded by the incorporation of 5 mg. of benzocaine per troche. The local anesthetic action so provided, in addition to the specific antibiotic effect of the bacitracin, makes for a wider field of usefulness in the treatment of pharyngitis, tonsillitis, and gingivitis. These troches remain intact for a prolonged period thereby exerting an extended anesthetic influence and creating high salivary bacitracin levels.

The confection-like chocolate taste of C.S.C. Bacitracin Troches with Benzocaine makes for universal patient acceptance. Their candy-like taste and appearance encourage their continued use by adults as well as by older children.

C. S. C. Bacitracin Troches with Benzocaine are available on prescription through all pharmacies in bottles of 25. C.S.C. Pharmaceuticals a division of commercial solvents corporation 17 EAST 42ND STREET, NEW YORK 17, N.Y.



Each C.S.C. Bacitracin Trocke with Benzocaine contains

## Forensic Medicine

ARTHUR L. H. STREET, LL.B.

Prepared especially for Modern Medicine

PROBLEM: In a murder trial, was an associate toxicologist in a state department of toxicology, who had had considerable experience in performing autopsies and spectroscopic and ballistic examinations, qualified to testify as to whether wounds on the victim's body were inflicted before or after death?

#### COURT'S ANSWER: Yes.

The Alabama Supreme Court mentioned the witness' explanation that an antemortem wound is indicated by a swelling of the area and interstitial hemorrhage (48 So. 2d 553).

PROBLEM: A Wisconsin statute, as in other states, forbids a physician to disclose, without a patient's or the patient's representative's consent, information acquired in a professional capacity to enable the doctor to treat the patient. In a third party's suit against a deceased patient's estate on a note, did the trial judge wrongfully refuse to permit the decedent's doctor, who witnessed signing of the note, to testify to the circumstances under which the note was signed?

#### COURT'S ANSWER: Yes.

The Wisconsin Supreme Court said that under the statute a physician "is a competent witness if his part in the transaction was disassociated from his professional duties or where the privilege with relation to confidential communications has been waived" by the patient or his representative (44 N.W. 2d 574).


PROBLEM: In a wife's suit to compel her husband to support her infant child, defended on the ground of nonpaternity, did the trial judge wrongfully reject the husband's request that the wife and child be required to take a blood-grouping test, when the wife gave no good reason for not submitting?

#### COURT'S ANSWER: Yes.

The Appellate Division of the New Jersey Superior Court cited numerous legal and medical authorities as vouching for the accuracy of such tests, "not to prove paternity, and not always to disprove it," but as disproving it "conclusively in a great many cases provided they are administered by specially qualified experts."

The court said that, although a trial judge has considerable discretion to order a blood-grouping test, his refusal to do so is subject to reversal on appeal when the question of parentage is relevant and a test may have crucial effect.

But the court recognized that refusal to require a blood test might be excused on proof that submission to it might endanger the patient's health. However, because the giving of a few drops of blood can rarely involve any risk, the person refusing to submit must establish the possibility of danger. Here the wife showed no medical reason for refusing.



## Relationship of Stress to Autonomic Lability

Studies have shown that functional disorders often are a result of the patient's inability to adjust to emotionally stressful

situations (stressor factors).

Nervous tension and chronic anxiety, discharged through a labile Autonomic Nervous System, can cause somatic disturbance. Such states may involve any one of the organ systems or several at one time. The outline below relates gastrointestinal and cardiovascular symptomatology to the exaggerated response of the autonomic nervous system.

	Physiologic Effects of Autonomic Disch			
	Sympathetic	Parasympathetic		
Gastro- intestinal	Hypomotility Intestinal Atony Hyposecretion Reduced salivation	Hypermotility Gastrointestinal spasm Hypersecretion		
Cardio- vascular	Rapid heart rate Peripheral vaso- constriction	Slow heart rate Vasodilatation		
Functional Manifesta- tions	Palpitation Tachycardia Elevated B. P. Dry mouth— throat	Heartburn Nausea-vomiting Low B. P. Colonic spasm		

Diagnosis of functional disorder is supported by the following indications of autonomic lability:

Variable Blood Pressure; Body Temperature Variations; Changing pulse rate; Deviations in B. M. R.; Exaggerated Cold Pressure Reflex; Glucose Tolerance Alterations.

Therapy in these cases is directed toward: 1) relief of symptoms by drug, therapy (so making the patient more amenable to psychotherapy); 2) psychotherapeutic guidance in making adjustment to stressful situations and correction of unhealthy attitudes.

Clinicians report that good therapeutic results are produced by combined adrenergic (ergotamine) and cholinergic blockade (Bellafoline) with central sedation (phenobarbital). A convenient preparation of this nature is available in the form of Bellergal Tablets. Full data on request; write to:

## Sandoz Pharmaceuticals

DIVISION OF SANDOZ CHEMICAL WORKS, INC. 68 CHARLTON STREET, NEW YORK 14, N. Y. The court assumed, without deciding, that a defendant in a criminal case might not be required to submit to a blood-grouping test, but decided that in civil cases "appropriate and familiar judicial sanctions" might be used to enforce obedience. This would seem to imply that a wife's suit to compel support of her child might be dismissed if she persisted in refusing to submit to the test (76 Atl. 2d 717).

PROBLEM: Kansas, as other states, has separate statutory provisions for the licensing of osteopaths, as distinguished from physicians and surgeons. Are the constitutional rights of osteopaths violated because they cannot practice medicine or surgery, although they may have been trained in those fields of practice in osteopathic schools?

#### COURT'S ANSWER: No.

In disposing of the question, the U.S. District Court for Kansas did not decide whether graduates of such schools could practice "incidental surgery such as severing the umbilical cord or repairing lacerations by . . . surgery in obstetrical cases."

The court noted that since 1913 osteopathic schools have broadened their curricula so as to teach limited drug therapy and surgery, but declared that they could not thereby broaden a license to practice osteopathy into a license to practice medicine and surgery in disregard of the statutes regulating that practice.

The court also recognized the reasonableness of exacting from osteopaths enough knowledge concerning medicine and surgery to help them recognize the need for referring patients to licensed doctors for treatment and surgery falling outside the practice of osteopathy (92 Fed. Supp. 280).



"Premarin" with Methyltestosterone wherever combined estrogen-androgen therapy is indicated.

In the treatment of fractures and osteoporosis in either sex to promote bone development, tissue growth, and repair.

# **e.g.**

"PREMARIN".

In the treatment of the female climacteric in certain selected cases.

"Premarin" with Methyltestosterone wherever combined estrogen-androgen therapy is indicated.

#### WITH

#### METHYLTESTOSTERONE

In the treatment of on the basis that a

In the treatment of dysmenorrhea in an attempt to suppress ovulation on the basis that anovulatory bleeding is usually painless.

for combined estrogen-androgen therapy

In the treatment of the male climacteric to reduce follicle-stimulating hormone levels.

"Premerin" with Methyltestosterone provides a steroid combination designed to permit utilization of both the complementary and the neutralizing effects of estragen and androgen when administered concomitantly. Thus certain properties of either sex hormone may be employed in the opposite sex with a minimum of side effects.

Availability: Each tablet provides estrogens in their naturally occurring, water-soluble, conjugated form expressed as sodium estrone sulfate, together with methyltestosterone.

No. 879-Conjugated estragens equine

AYERST, McKENNA & HARRISON LIMITED
22 East 40th Street,
New York 16, New York



	("Premarin")	1.25	mg
	Methyltestosterone	10.0	mg
	Bottles of 100 tablets (yellow)		
No. 87	-Conjugated estragens equine		
	("Premarin")	0.625	mg.
	Methyltestosterone	5.0	mg
	Bottles of 100 toblets (red)		



Each tablet contains:

Veratrum viride 100 mg. Mannital hexanitrate 15 gr. Rutin

> Phenobarbital % or.

10 mg.

**VERUTAL Tablets (RAND)** 

CONTAIN Veretrum

VIRIDE plus other

ACTIVE AGENTS. NO

SINGLE DRUG IS SUF-

FICIENT FOR THE COM-

PLETE TREATMENT OF

THIS COMPLEX DISEASE.

for effective treatment of HYPERTENSION

Clinical trial package and literature on request



pharmaceutical co., inc.

albany, n. y.

PROBLEM: A surgeon assisted a hospital nurse in moving an unconscious patient from an operating room on a wheeled stretcher. An elevator door was open and, while the doctor was attempting to bring the elevator to the floor level, the nurse carelessly permitted the stretcher to roll into the elevator shaft, injuring the patient. [1] Was the doctor liable in damages? [2] Was his partner, who was not present, liable?

#### COURT'S ANSWERS: Yes.

1 In this case, decided by the Iowa Supreme Court some years ago, the court does not seem to have adopted a view that a surgeon is under any implied responsibility for the safety of a patient in transit between his room and the operating room. The surgeon assumed responsibility in this instance when he undertook to help move the patient. Liability was imposed not on a theory that the doctor was liable for the nurse's carelessness, but because he failed to safeguard the patient when he knew that the nurse was about to release her hold on the stretcher to help him bring the elevator to the floor level.

The dissenting judge said that the decision greatly increased "the liability, already sufficiently burdensome, of the medical profession." He thought that the hospital, rather than the doctor, ought to be charged with the nurse's neglect, which was the direct cause of the accident.

2) Finding from the evidence that the surgeon had undertaken, on behalf of his firm, to assist in the removal of the patient from the operating room to her hospital room, the Supreme Court decided that the doctor's partner was automatically liable jointly with the doctor. The dissenting judge thought that the assistance given by the surgeon ought



When pregnancy is first diagnosed, the need for increased amounts of calcium, phosphorous, iron and vitamins is already present.

OBron-specifically designed for the OB patient-provides balanced proportions of calcium, phosphorous, iron and vitamins to meet the added nutritional demand of the mother and to safeguard the optimal development and growth of the fetus.

Especially beneficial during the period of lactation, OBron supplies adequate vitamins and minerals to protect the nutritional state of the mother and insure an optimal content of these nutrients in the milk for the nursing child.



## OBRON ...


#### CALCIUM · IRON · PHOSPHORUS · VITAMINS ... ALL IN ONE CAPSULE

*Dicalcium Phosphate, Anhydrous 768 mg.	Vitamin B <sub>2</sub> (Riboflavin) 2 mg.
Ferrous Sulphate U.S.P 64.8 mg.	Vitamin B <sub>6</sub> (Pyridoxine Hydrochloride) 0.5 mg.
Vitamin A (Fish-Liver Oil) 5,000 U.S.P. Units	Vitamin C (Ascorbic Acid) 37.5 mg.
Vitamin D (Irradiated Ergosterol) 400 U.S.P. Units	Niacinamide 20.0 mg.
Vitamin B <sub>1</sub> (Thiamine Hydrochloride). 2 mg.	Calcium Pantothenate 3.0 mg.

\*Equivalent to 15 grains Dicalcium Phosphate Dihydrate



536 N. Lake Shore Drive . Chicago 11, III.



essential hypertension

Maxitate with Rhamnotin and Maxitate with Rhamnobarb are ideal for routine treatment and protection because they:

- STABILIZE blood pressure.
- RESTORE and maintain vascular integrity and permeability.
- ELIMINATE exaggerated reactions to emotional responses.
- ARREST progression of the arteriosclerotic process.
- GUARD against the occurrence of cerebral vascular accidents.
- HAVE no known contraindications.
- · ARE SAFE in use.

description: Each Maxitate with Rhamnotin tablet green; contains \*Maxitate 30 mg., Rutin 15 mg., and Ascorbic Acid 20 mg. Each Maxitate with Rhamnobarb tablet (orange) contains \*Maxitate 30 mg., Rutin 15 mg., Ascorbic Acid 20 mg., and Phenobarbital 15 mg.

dosage: 1 to 2 tablets every 4 to 6 hours according to individual requirements.

\*The STABILIZED form of Mannitol Hexanitrate pioneered by Strasenburgh research.



Supply for amhating treatment available. Write. R. J. Strasenburgh Co. Dept. M. Rochester 14 N.Y.

to have been regarded as his voluntary and personal act and therefore not chargeable to the firm (115 N.W. 921).

In several states, appellate courts have applied to medical partner-ships the rule commonly applied to all business partnerships—that each member of a firm is its agent and guarantees to the public that each will use due care and skill to avoid injuring those who deal with the firm. But, of course, if one member of a firm injures an outsider in committing some act that falls outside the scope of the firm business, he alone can be held liable.

PROBLEM: Does a doctor's refusal to treat a Negro fall within a statute making it unlawful for the proprietor, keeper, or manager of a restaurant, barber shop, conveyance, theater, store, "or any other place of public accommodation or amusement" to deny to a citizen, except for reasons applicable alike to all citizens regardless of color or race, the "full enjoyment of the accommodations, advantages, facilities, or privileges thereof"?

#### COURT'S ANSWER: No.

The Ohio Court of Common Pleas for Montgomery County dismissed a suit for damages brought against a dentist under the statute. But the opinion treated physicians, surgeons, and lawyers, as well as dentists, as falling outside the law.

The court cited an opinion of an Ohio appellate court to the effect that a doctor is not liable for refusing to respond to any call unless he has contracted to render services.

The court said that it would be unreasonable to assume that the legislature intended, without expressly saying so, to require a doctor to

# When the diagnosis is Pyelitis



o establish and maintain urinary antisepsis, MANDELAMINE\* is many times preferred because it is quickly effective against the organisms most commonly encountered in urinary-tract infections. Its exceptional freedom from untoward reactions and its wide range of antibacterial activity commend it for use as soon as the diagnosis has been made.

Urinary antisepsis is often achieved in uncomplicated pyelitis in as few as three days. Speedy recovery is thus secured in many cases without necessitating highercost therapy. Renal insufficiency is the only major contraindication to MANDELAMINE therapy.

MANDELAMINE is available in bottles of 120, 500, and 1,000 enteric-coated tablets, through all prescription pharmacies. Comprehensive literature and samples for clinical trial will be furnished to physicians on request.



NEPERA CHEMICAL CO., INC.

Pharmaceutical Manufacturers. NEPERA PARK, YONKERS 2, N. Y.

<sup>\*</sup>MANDELAMINE is the registered trademark of Nepera Chemical Co., Inc., for its brand of methenamine mandelate,



## for Coughs...

in acute and chronic bronchitis and paroxysms of bronchial asthma . . . whooping cough, dry catarrhal coughs and smoker's cough—

## **PERTUSSIN**

with no undesirable side effects for the patient, helps Nature relieve coughs when not due to organic disease.

Its active ingredient, Extract of Thyme (Taeschner Process), acts as an expectorant and antispasmodic. It increases natural secretions to soothe dry, irritated membranes. It may be prescribed for children and adults. Pleasant to take.

Trial packages on request.

SEECK & KADE, INC. New York 13, N. Y. treat every prospective patient regardless of the doctor's special line. The court thought that having specially mentioned barbers, and so on, the legislature would have specified dentists, physicians, and surgeons had it been intended to include them in the statute (95 N.E. 2d 30).

PROBLEMS: An orderly in a physician's private hospital assisted in childbirth when the doctor was away and no other doctor could be secured. Did the orderly violate a statute forbidding practice of medicine without a license but excepting "the furnishing of medical assistance in a case of emergency"? The orderly had been a surgical technician in the Medical Corps. Was he guilty of violating a statute forbidding an unlicensed person to hold himself out as being able to diagnose and treat human disease or to operate because he was commonly known as "Doctor B"?

#### COURT'S ANSWERS: No.

The Florida Supreme Court said that the man exceeded his authority as nurse and orderly by assisting in delivery of a child, but that the emergency justified what he did. Apparently the statute did not specifically ban use of the word "doctor," and the court noted that the prefix "Doctor" is commonly applied by the public to druggists, nurses, and others (47 So. 2d 764).

PROBLEM: A malpractice suit was decided in a doctor's favor upon the ground that the plaintiff, a woman, failed to substantiate a claim that her child was delivered dead through the doctor's refusal to attend her. The husband was a material witness as to the doctor's refusal to continue attendance. The husband testified that the doctor had been paid in full for all services to date when allegedly he refused to

(Continued on page 46)



Your Surgical Dealer Wants To Show You



## THIS COMPACT HANDFUL

Here's an especially practical diagnostic set. Two fine, dependable instruments in a scuff-proof, stain-proof, washable zipper pouch of smooth Vynyl . . . that really fits within the pocket.\*

Or, if preferred, one instrument, Otoscope, pouch contained. Attractively priced, too.

Ophthalmoscope details: National's No. 40 with one finger control for apertures, dioptres, light and with brilliant light beam and three epertures plus red-free filter; dioptric range -20 to +25. With smooth precision action and precision vision. Head . . . metal, wafer-thin, light, black.

Otoscope details: National's No. 22 with perfected, brighter illumination and traditional money-saving flashlight bulb. Head housing . . metal, black. Positive, no fuss, lazy-latch speculum holder. In-vision parts—black . . no distractions. Specula holder provides rectilinear adjustment. Head and holder . . all-metal. Impact-proof, guaranteed specula, six, in graduated sizes.

\* Both are new, improved instruments of their kind . . , lifetime guaranteed.

A valuable addition to your reference library — write for it today: "What The General Practitioner Should Know About Ophthalmoscopic Examinations."

Use coupon below.

n	A	TI	0	n	Ð	
			U RUMEN			
			., Elmhu			

World-Famous Makers of Electro-Medical Instruments

-	_	_									
M.	A: -	1	Plant.	1			4	0-	1	~	١
Na	шо	naı	Electr	IC I	nsu	4111	ent	LO.,	Inc	. 4	,
n.	-	2424	92-21	Car	-	Awa	FI	mburo			
v	EDI.	100 M	37.71	-	una	MAG		minut 9			

- Send me my free copy of "What The General Practitioner Should Know About Ophthalmoscopic Examinations."
- Send me additional data on the National "Pocket" Otos.-Ophth. Set

Nome ......

City\_\_\_\_\_Zone\_\_State\_\_\_

ta



Winthrop Stearns ....

New York 18, N. Y. Windsor, Ont.

# re you puzzled...

## in choosing a bulk laxative?


...then try Mucilose-highly purified hemicellulose of Plantago loeflingii. Mucilose reeducates the constipated intestine to "physiologic elimination" by providing soft, demulcent bulk and by restoring the intra-intestinal "water balance." In addition, Mucilose is hypoallergenic and does not interfere with digestion.

Obtainable in 4 ounce and 16 ounce containers in the following forms:



Mucilose Flakes Concentrated Mucilose Flakes (special formula with dextrose) Mucilose Granules (special formula with dextrose) Mucilose with Cascara Granules

# UCILOSE®



for physiologic elimination



continue. The doctor testified that the husband discharged him and that his fees were not paid. Under these circumstances did the trial judge properly permit the doctor to show by the operator of a credit bureau that the husband told the latter that if the doctor should sue him on the account he would sue the doctor for malpractice?

#### COURT'S ANSWER: Yes.

This case was decided by the Oklahoma Supreme Court (215 Pac. 2d 827).

PROBLEM: Preparatory to a fistula operation and before the patient's surgeon arrived, a private hospital attendant burned the patient, who was under anesthesia, by negligently applying to the patient's body an electric plate to which was attached an electrocautery knife. [1] Was the doctor liable? [2] Was the hospital liable?


COURT'S ANSWERS: [1] No. [2] Yes.

The New York Supreme Court, Brooklyn, said that suit for damages against the surgeon was properly dismissed because the jury had specifically decided that, under the circumstances, it was not necessary for the doctor to supervise the placing of the plate.

As to the liability of the private hospital operated for profit, the court pointed out that another New York court had decided that such a hospital was liable for negligence of an intern and that, for stronger reasons, there should be liability for "carelessness of a nurse or other agent or employee of a private hospital." The court distinguished the case from one in which it was decided that a public or charitable hospital was not liable for negligence of a nurse while performing professional duty (99 N.Y. Supp. 2d 814).

# FREE TO PHYSICIANS...

## New Low-Sodium Reducing Diets



The value of sodium-restricted diets in congestive heart failure, high blood pressure, and certain forms of renal disease has been established by numerous clinical studies. In patients requiring a reduction of weight with a sodium-restricted diet, specification of **DIETENE®** provides a truly palatable means of accomplishing the desired result without sacrificing nutritional adequacy, or jeopardizing the low-sodium regimen.

**DIETENE** is an excellent low-calorie source of biologically superior proteins plus protective amounts of essential vitamins and minerals.

SUPPLIED: In 1-lb. cans, plain or chocolate flavor, available through all pharmacies at \$1.55, PATIENTS ENJOY TAKING DIETENE.

SEND FOR YOUR FREE SUPPLY OF 1000-CALORIE, LOW-





Not advertised to the laity

THE DIETENE COMPAN
--------------------

Dept. DM 21

518 FIFTH AVENUE SOUTH, MINNEAPOLIS 15, MINNESOTA

Please send me a free supply of the new 1000-calorie DIETENE Reducing Diet, providing only 0.7 to 1 Gm. of sodium per day.

City.......State.....

## Washington Letter

#### Effective Mass Treatment Is Goal of Civil Defense Manual

Many hours of behind-the-scenes discussion went into preparation of the statement on treatment of burns included in Civil Defense Administration's health services handbook, now being circulated throughout the country to medical leaders. The problem facing National Security Resources Board's staff was to reconcile the many burn treatments currently in use, but limit the medical material in stockpiles.

Representatives of several government departments and professional associations were consulted on this, as they were on other controversial points.

The method selected is a dry treatment, employing a cellulose pad faced with extremely fine gauze and backed with coarse gauze, held in place with tensile roller bandages.

The treatment was not selected as

(Continued on page 52)



## new clinical studies'

again prove value of

Westhiazole Vaginal in cervicitis and

vaginitis. Useful in clearing up cervical mucous

plug or mucopurulent discharge; promotes

"rapid healing" after cauterization; "gratifying results"

when applied before and after hysterectomies and plastic repair.

### westhiazole vaginal

dainty, convenient single-dose disposable applicators

send for samples and reprint 1

by Stein, I. F. and Kaye, B. M.: Su. Clin. North Am. 30:259, 1950.

WESTWOOD PHARMACEUTICALS
Division of Foster-Milburn Co.
468 Dewitt St., Buffalo 13, N. Y.

WESTHIAZOLE VAGINAL:
a sterile jelly,
10% SULFATHIAZOLE,
4% UREA, 3% LACTIC
ACID, 1% ACETIC ACID
in a polyethylene
glycol base. Acidifies,
combats secondary
infection, speeds healing.



# Cortone



Saline Suspension of Cortone Acetate

(1 cc.=25 mg.) vials, 20 cc.

Clinical studies have demonstrated that the therapeutic activity of Cortone\* is similar whether administered parenterally or orally. Dosage requirements are approximately the same, and the two routes of administration may be used interchangeably or additively at any time during treatment.

Although the manufacture of Cortone—probably the most intricate and lengthy synthesis ever undertaken—has imposed unprecedented difficulties, every effort is being made to increase production and, in the meantime, to achieve an equitable national distribution of this vital drug.

Literature on Request

Key to a New Era in Medical Science



(CORTISONE Acetate Merck)

(11-Dehydro-17-hydroxycorticosterone-21-acetate)

Among the conditions in which Cortone has produced striking clinical improvement are:

RHEUMATOID ARTHRITIS and Related Rheumatic Diseases

ACUTE RHEUMATIC FEVER

ALLERGIC DISORDERS, including Bronchial Asthma

INFLAMMATORY EYE DISEASES

SKIN DISORDERS, notably Atopic Dermatitis, Psoriasis, Exfoliative Dermatitis, including cases secondary to drug reactions, and Pemphigus

LUPUS ERYTHEMATOSUS (Early)

ADDISON'S DISEASE

\*CORTONE is the registered trade-mark of Merck & Co., Inc. for its brand of cortisone.



MERCK & CO., INC.

Manufacturing Chemists

# What is

# GANTRISIN

A new, safer sulfonamide with a wider antibacterial spectrum.

#### same indications as other sulfonamides?

More; it has been effective in some infections

not responsive to other sulfonamides and antibiotics.

#### how about toxicity?

High solubility prevents renal blocking. Incidence of other reactions is also very low.

# 'ROCHE'?

#### should the patient be alkalized?

Not necessary with Gantrisin® because of its high solubility.

#### how about cost?

Gantrisin is so economical that it can be prescribed without straining the patient's budget.

#### HOFFMANN-LA ROCHE INC.

Roche Park • Nutley 10 • New Jersey

# LOWEST NICOTINE BY FAR



#### Nicotine Actually Bred Out Of The Leaf

The low nicotine content of the tobacco in John Alden eigarettes has been achieved by painstaking tobacco-plant breeding over a long period of years. Tests\* indicate 85% less nicotine in the moke of John Alden eigarettes than in four leading popular brands tested—75% less than in two leading denicotinized brands tested.

#### **Importance To Doctors And Patients**

John Alden eigarettes ofter a satisfactory solution to the problem of reducing a patient's nicotine intake. They accomplish the reduction often necessary for a patient's improved physiological condition without imposing on the patient the strain of breaking a pleasurable habit. At the same time they free the doctor from the distasteful task of prescribing "No smoking" and assure him that the possibility of nicotine contributing to or aggravating the patient's symptoms has been reduced to a minimum.

### ABOUT THE NEW TOBACCO

OWEST " NICOTINE " IM

John Alden cigarettes are made from a completely new variety of tobacco. This variety was developed after 15 years of research by the Kentucky Agricultural Experiment Station. Because of its extremely low nicotine content, it has been given a separate classification, 31V, by the U. S. Department of Agriculture.

#### Also Available: John Alden Cigars and Pipe Tobacco

\*A comprehensive series of smoke tests conducted by one of the country's leading independent consulting laboratories. A summary of these results is available on request.

## FREE PROFESSIONAL SAMPLES

11 West 42nd St., N. Y. 18, N. Y. Dept. M2		
Send me free samples of John Alden Cigarettes.		
Name	M	
Address		

the ideal therapy, NSRB was careful to point out, but because it met the following requirements:

1] The procedure is effective and provides relief from pain.

2] It is applicable to all, or nearly all, types of burns, and can be employed by first-aid workers.

3] It is usable under most field conditions, is suitable for rapid ap-



Applying cellulose pad

plication, and requires short time for application and after care.

4] Necessary supplies can be stockpiled and are relatively inexpensive.

5] The method does not predetermine subsequent treatment.

The handbook is modest in its description of the method selected. The report says:

A large variety of treatments for burns currently is in use. The multipli-



# Quality and Craftsmanship ...



HOW AVAILABLE: GELUSIL\* Warner, the safe, effective and reliable antacid preparation is purely local and non-systemic in its action.

TABLETS—each containing magnesium trisilicate, 0.5 Gm (7.5 grains) and dried aluminum hydroxide gel, 0.25 Gm (4 grains): boxes of 50 and 100, and bottles of 1000 tablets.

LIQUID—magnesium trisilicate, 0.5 Gm (7.5 grains) and aluminum hydroxide, 0.25 Gm (4 grains) per 4 cc (1 teaspoonful): bortles of 6 and 12 fluidounces.

Seley, S. A.: Medical Management of Pyloric Obstruction Resulting from Peptic Ulcer, Am. J. Dig. Dis., 13:238, 1946.

\*T. M. Reg. U. S. Pat. Off.

# GELUSIL Warner

Once in a long while a remedy is evolved which meets practically all of the medical requisites: effective, safe, and reliable. In the management of peptic ulcer or hyperacidic conditions, GELUSIL® "Warner" by combining comparatively non-reactive aluminum hydroxide gel with magnesium trisilicate, provides the advantages of both.

Prompt action Prompt relief
Prolonged action Prolonged relief

without secondary acid rise, chloride depletion, or danger of alkalosis; and, most important, there is practically no conscipation.<sup>1</sup>

WILLIAM R. WARNER

Division of Warner-Hudnut, Inc. New York • Los Angeles • St. Louis city of methods indicates that none is strikingly superior to the others and that much more research work in the whole field is needed. The body area involved, the degree and cause of the burn and other variable factors may demand some variations in any kind of treatment developed. Each physician should be free to use the type of external treatment which his judgment dictates, providing that supplies and personnel for the type of treatment are available. . . .

The criteria for a satisfactory local burn treatment for the thousands of burn casualties that would result from atomic attack are considerably different from those which would be used to evaluate an individualized type of local treatment. In the latter case, one would search for an ideal treatment for a particular type of burn. With mass casualties, however, the need is for an effective method which could be used for all cases and which would be adaptable to mass-treatment methods.

The dry treatment, NSRB says, "meets the criteria better than any other existing method. It is a relatively new regimen, based on the principle of a dry dressing. It fur-

nishes a covering which relieves pain and bars further introduction of pathogenic organisms. Its absorbability will prevent rapid total saturation with exudate which would lead to the necessity for undesirable frequent changes of dressing."

Civil Defense Administration, which came into operation just in time to issue the book after it had been prepared by

NSRB, is anxious to get a copy into the hands of every physician and every civil defense worker as rapidly as possible.

The handbook may be ordered by title, Health Services and Special Weapons Defense, from Superintendent of Documents, Government Printing Office, Washington 25, D.C. The price is 60¢ a single copy, with a 25% discount in lots of 100.

Every conceivable health phase of civil defense preparation and operation is dealt with, from hospital expansion to morgue identification. Physicians—who will be responsible for detecting and identifying bacteria or gases which may possibly be used—should be particularly interested in chapters on radiologic and biologic defense. Other fields covered include: training and first aid, expansion and improvisation of hospitals, health supplies, water and other sanitation services, laboratory,

(Continued on page 57)



"Just say we're sorry to decline her kind invitation but that I'll gladly hear her symptoms any day between two and four."

FOR THE CONSTIPATED PATIENT...

# Laxative (ACTION WITHOUT REACTION



Phospha-Sada (Fleet) he long been authoritativel recognized for its dependable efficacy and desirable qualitie in the treatment of intestination stasis. In average doses, produces a soft and formed, rathe than a watery, evacuation; and its gentle action is quite free from irritation, griping, early tendency toward habituation, or other adverse reaction.


Phospho Soda Fleet is a solution containing in each 100 cc sod. In phosphote 48 Gm and sodium phosphote 18 Gm Both Phospho Soand Fleet are resolved trademarks. 1 ( 8 Fleet Co.

C. B. FLEET COMPANY, INC., LYNCHBURG. VA

THERE IS ONLY ONE

# PHOSPHO-SODA (FLEET)

A Laxative for Judicious Therapy



# in biliary tract disorders

Hydrocholeresis with *Decholin* and *Decholin Sodium* produces a gentle lavage of the biliary tree. Copious, fluid bile flushes away mucus, pus and thickened bile and re-establishes normal diagrams.

# for best results

Hydrocholeretic therapy should be extended through the optimal treatment period. An average dose of Decholin is 1 or 2 tablets three times daily for four to six weeks. Prescription of 100 tablets is recommended for maximum efficacy and economy. The course may be repeated after an interval of one or two weeks if desired. For more rapid and intensive hydrocholeresis, therapy may be initiated with Decholin Sodium.

# DECHOLIN

Decholin tablets (brand of dehydrocholic acid) of 3¼ gr. (0.25 Gm.), in bottles of 100, 500, 1,000 and 5,000.

Decholin Sodium (brand of sodium dehydrocholate) is supplied in a 20% solution for intravenous administration. 3 cc., 5 cc. and 10 cc. ampuls — boxes of 3, 20 and 100.

Decholin and Decholin Sodium, trademarks reg.

AMES COMPANY, INC . ELKHART, INDIANA

Ames Company of Canada, Ltd., Toronto



nutrition, industrial, and veterinary services, and vital records and reports.

While the handbook is only advisory, in the face of enemy attack Civil Defense Administration could be authorized by the President to require communities and individuals to carry out the recommendations.

#### Nurse Supply Critical

By July 1, all three military services expect to have at least doubled the number of nurses they had on active duty at the first of the year. Some will come from reserves, but the vast majority will be recruited from military-age civilian nurses who have not had previous military duty.

Prospects are that at least 10,000 nurses will be withdrawn from civilian life. This, according to the American Nurses' Association, should more than equal the number of nurses to be graduated this year. Furthermore, Red Cross, Public Health Service, and new civil defense facilities will require many thousands of additional nurses.

Defense planners are attempting to meet both the immediate and the long range problem. One temporary approach, already begun, is the mass training of nurses' aides, with the prospect that many thousands will be ready for service by summer. Operating the way they did in World War II, nurses' aides will permit the supply of registered nurses to be spread thinner as demands increase.

For permanent solution, nurses' organizations and defense officials are prepared to support legislation for federal aid to nursing schools and teaching hospitals. The plan as now described will be a liberal ver-

# From where I sit



Watch Out For The "Blind Spots"

Stopped by Squint Miller's farm the other day and saw a vinegar bottle in his kitchen with an oversized cucumber inside it. The cucumber filled the whole bottle.

"What's a cucumber doing in' there?" I asked him. "That's my 'blind spot' reminder," says Squint. "My grandmother kept one in her kitchen to remind her to take stock of herself now and then.

"I slipped that bottle over the cucumber when it was just starting to grow on the vine," he went on. "And like certain viewpoints, not noticed, it just grew and grew—now it's there to stay."

From where I sit, we could all take a cue from Squint and watch out for our own "blind spots." Sometimes we impose our views on our neighbor without thinking of his rights as an American—his right to follow his profession where and how he chooses, or say, his right to enjoy a glass of beer now and then. We won't be tripped by "blind spots" if we keep our eyes—and minds—open!

Joe Marsh

Copyright, 1951, United States Brewers Foundation



# Dioloxol

TABLETS · CAPSULES · ELIXIR

The new muscle relaxant for certain spastic and neuromuscular conditions which acts through the brain and spinal cord

"...in an unusual pharmacologic niche."



parkinson's syndrome hemiplegia diplegia infantile cerebral palsy tetanus low back pain other muscular spastic disorders anesthesia

Descriptive literature and specimens available

G. W. Carnrick Co.

NEWARK 1, NEW JERSEY

9 MJ9

sion of the nurses' section of the aid to medical education bill, which was allowed to die in the last Congress.

Concurrently, American Nurses' Association is leading two campaigns, one to induce nurses to volunteer for Army duty, the other to replenish the civilian supply by persuading young women to take nurses' training. The association is also urging retired nurses to return to duty, thereby making possible the release of military-age nurses for military service.

#### Local Public Health Bill

Because of the national emergency, the new bill by Rep. Percy Priest (D., Tenn.) for federal aid to local public health units has a good chance of passage at this session of Congress. The proposal is a revamped version of other bills on the subject, with emphasis on defense areas and a new definition of public health functions, modified to meet criticism of earlier definitions. Authorizations would cease when the current national emergency is declared ended: a state would give priority to defense areas and could start its plan before all sections were included; and the surgeon general would define defense areas, after consulting with other government officials.

States would be allowed various amounts of money, depending on their population and per capita income, all of which would be passed on to the local units. In no case could the federal contribution exceed \$2 per capita.

#### Washington Notes

"YOUR BEST BUY," a competently prepared pamphlet designed to sell





# Announces A new Multivitamin Capsule for patients of all ages



Small, attractive, easy-to-swallow, Mead's POLY-VI-CAPS® contain six essential vitamins in well-balanced amounts. Children especially will like these bright black and orange two-color capsules.

POLY-VI-CAPS meet physicians' requirements for a small multivitamin capsule with unusual versatility of use. Dosage can be adjusted easily to suit the varying needs of the individual patient. POLY-VI-CAPS are economically priced too! Available in bottles of 100 capsules.

#### Each Poly-Vi-Cap supplies:

Vitamin A	3000 U.S.P. units
Vitamin D	400 U.S.P. units
Thiamine	1.0 mg.
Riboflavin	1.2 mg.
Niacinamide	8.0 mg.
Ascorbic Acid	50.0 mg.

SPECIFY



MEAD JOHNSON & CO



# Quick bite — up all night

The "eat and run" type patient often pays the penalty for haste with discomfort from hyperacidity. A good way to provide fast, effective relief is to recommend BiSoDoL. This modern, dependable antacid formula acts quickly and sustains relief for a long period of time. BiSoDoL has a pleasant taste and is well tolerated. For an efficient antacidrecommend

BiSoDoL\*

tablets or powder

WHITEHALL PHARMACAL COMPANY 22 East 40th Street, New York 16, N.Y. the country on the advantages of local public health services, was released by Public Health Service at just about the time Mr. Priest introduced his new bill. It undoubtedly will be effective in building up back-home support.

APPOINTMENT of highly competent Rear Adm. H. Lemont Pugh as surgeon general by President Truman was in some respects a surprise. The deputy (Adm. Pugh held the post four years) usually is not elevated.

with \$3,000,000 from Defense Department, and another \$9,000,000 earmarked, Red Cross is able to launch properly a tremendously complicated blood collection campaign. The problem now is to find professional personnel, without unduly interfering with the military services, public health and the Veterans Administration.

JOHN L. THURSTON, who has done a good job for FSA Administrator Oscar Ewing for several years without a title, now is officially Ewing's deputy, with title of assistant.

In CHALLENGING Mr. Ewing on federal regulations governing assistance to the needy, Gov. Dewey has uncarthed an old and never-settled issue: At what point do federal regulations cease to protect federal money and begin to encroach on state prerogatives? The question is raised at one time or another regarding almost every grants program—with a different answer each time.

CHAIRMAN VINSON of House Armed Services Committee indicated he might favor an Air Force Medical Center if proposed in a separate bill. His committee killed the plan when it was a part of another bill.

# for estrogen therapy PROGYNON\*\* the true, natural

estrogenic hormone

When the menopause necessitates substitution therapy, quicker, more certain benefit without side effects will be obtained by administering and prescribing the Progreson preparations of estradiol, the hormone elaborated by the human ovary.



PROGYNON-B®

PROGYNON buccal tablets

PROGYNON-DP®

PROGYNON, MICROPELLETS®

PROGYNON pellets

PROGYNON tablets

PROGYNON-DH® ointment

**PROGYNON-DH** suppositories


Schering corporation-bloomfield, New Jersey

PROGYNON





"Sorry sir, but the diathermy treatment will have to wait until after Howdy-Doody."



# **g**lycerite

of Hydrogen Peroxide ife with Carbamide

Instill one-half dropperful into affected ear four times daily Supplied in one-ounce bottles with dropper

Samples and Literature on request

Constituents: Hydrogen Peroxide 1 5% Urea (Carbamide) 2.5% 8 Hydroxyquinoline 0 1%

Dissolved and stabilized In substantially anhydrous glycerol q s ad. 30cc.

International Pharmaceutical Corporation

132 Newbury Street, Boston 16, Massachusetts



The skill of the modern physician and surgeon is one of the minutes of our Yest, even a medical genius would be handicapped if he couldn't depend implicitly a shore to whom he must leak for his tools—the diagnostic medic, the countestes, and drugs with which he has learned as relieve pair, conquer illness and turn back do. We who supply the profession have no margin for error, no room for compose with scientific and business integrity. Mallinchrode has recognized this for Eq yes since the day in 1867 when Gustavus, Edward and Otto Mallinchrode founds company dedicated to the production of medicinal chemicals of the highest put and assured efficacy.

We are proud that these generations of medicinal and surgical practitioners and species have regarded Mallinchrode on a important ally, an operated accessary to their the

have presentions of motival and surgical practitioners and special-linel root or an important ally...an countrial accounty to their skill.

MALLINCREORY CREMICAL WORES . St. Louis . Now York



DO - ENPPURAN - STHEE FOR STREETA - PROCEPURATED BAR POLIATE - SO DA LIME in 1500 Phoesistical Chimicals

Wyeth

sofe . . . Substantially free

of untoward byeffects related to blood pressure, heart rate, or

central nervous stimulation.



## LETS THE PATIENT SLEEP AT NIGHT EFFICIENT NASAL DECONGESTANT...

Relief begins in minutes—lasts for hours.

Because WYAMINE is notably free of undesirable side-actions common to other vasoconstrictors, it is SAFER to use in individuals suffering from high blood pressure, heart dis-

ease, diabetes, or thyroid disease.

Available as: SOLUTION WYAMINE SULFATE, Bottles of 1 fl. oz.

WYAMINE-TYROTHRICIN NASAL SOLUTION,
Bottles of 1 fl. oz.

Wyamine-Penicillin, Capsules, Penicillin with Vasoconstrictor, for preparation of nasal solutions.

# WYAMINE

N-methylphenyl—tertiary—butylamine WYETH
WYETH INCORPORATED

Philadelphia 2, Pa.

#### MODERN MEDICINE

### Clinical Physiology of Potassium

WILLIAM S. HOFFMAN, M.D.\* University of Illinois, Chicago

THE human body contains about 160 gm. of potassium, most of which is in tissues, with approximately 2.7 gm. in extracellular fluids.

This unstable balance is probably maintained by the oxidative energy

of cells. Relations are constantly upset because serum levels change much faster than cellular concentrations can be shifted through slowly permeable membranes. Radioactive potassium injected intravenously takes fifteen hours to reach an equilibrium with total body concentration.

The average serum potassium for healthy subjects is 4 to 5.4 mEq. per liter, or 16 to 21 mg. per 100 cc.

Factors tending to raise serum levels are [1] increased potassium intake, [2] oliguria or anuria, [3] tissue breakdown or anoxia, [4] dehydration, and [5] adrenal cortical deficiency.

Serum values are lowered by [1] limited food intake, [2] increased renal excretion through diuresis, acidosis, or adrenal cortical hyperactivity, [3] dilution with potassiumfree fluids, [4] vomiting or diarrhea, and [5] glucose uptake by cells.

& Clinical physiology of potassium, J.A.M.A. 144:1157-1162, 1950.

A healthy adult usually ingests 70 to 100 mEq. of potassium daily, but stores practically none, explains William S. Hoffman, M.D. The renal mechanism is designed to remove a surplus rather than to conserve a deficit.


Gastrointestinal fluids contain high concentrations, the greater part normally absorbed back into the blood. With vomiting or diarrhea, 10 to 20 times the amount ordinarily excreted in feces may be lost, often when intake is poor.

For every gram of nitrogen broken down in destroyed cells, about 2.4 gm. of potassium is released. During starvation, trauma,

or disease, the substance may be added to serum faster than the removal rate.

Glycogenesis transports potassium from extracellular fluid into liver and muscle, causing a transient depletion of serum, and the fall may be increased by glucose ingestion, insulin, or epinephrine. If serum concentration is already low, serious symptoms may result, as in familial periodic paralysis.

When potassium is excreted by kidneys or bowel, much is accom-



Electrocardiographic changes produced by changes in serum potassium concentration

Modern Medicine, Feb. 15, 1951

panied by extracellular chloride. Since the loss in anions is made up by rise in bicarbonate, acute potassium deficiency is generally associated with alkalosis. The condition will be exacerbated by administration of sodium chloride and reversed by potassium chloride.

Low serum potassium nearly always means potassium deficiency, though rarely of the corresponding degree. But a deficit may be associated with normal or high serum values, probably owing to tissue breakdown.

The symptoms of hyperpotassemia chiefly involve the heart. The T wave of the electrocardiogram becomes taller, the QRS complex broader, and the P wave may disappear. Since potassium intoxication may occur with severe cellular depletion, replacement must be slow and preferably by stomach.

Low serum potassium broadens the T wave, lengthens the Q-T interval, and depresses the S-T segment. Symptoms include weakness, anorexia, lethargy, bowel distention, and lung edema.

Diabetic acidosis and polyuria result in large potassium losses from serum and body cells. If the patient is seriously dehydrated and acidotic, doses of insulin, isotonic sodium chloride, and sodium bicarbonate or lactate expand extracellular fluid and thus reduce the potassium level. Improved renal function and glycogenesis may then cause further and possibly fatal depletion.

Persons in deep diabetic coma should have potassium chloride by mouth or stomach tube in 2-gm. doses. A liter of intravenous fluid should not contain more than 2 gm. of potassium chloride.

For infantile diarrhea, isotonic sodium chloride solution and 5% dextrose are given with blood or plasma. When circulatory and renal functions are restored, Darrow's solution, containing 0.4 gm. of sodium chloride, 0.26 gm. of potassium chloride, and 0.59 gm. of sodium lactate, is injected subcutaneously, or diluted for oral therapy.

To correct surgical potassium deficit from tissue injury and alarm reaction, 2 gm. of potassium daily is added to the usual alimentation with saline solutions, dextrose, and amino acid solutions.

Potassium levels with *uremia* may be low, normal, or high, depending on type of renal damage, gastrointestinal reactions, and other factors. Medication containing potassium, often used for uremic acidosis, may be dangerous.

In states of dehydration, intravenous infusions should first restore blood volume, oxygen-carrying power, blood pressure, renal circulation, and acid-base balance. Ringer's solution may be employed for the initial injections. With improvement, potassium may then be started by mouth or, if necessary, by vein.

No extensive intravenous potassium management should be undertaken unless adequate laboratory facilities are available for quick potassium determinations. In fact, no program of repair treatment is safe without frequent determinations of urinary volume, specific gravity, pH, serum chloride, bicarbonate, sodium, and potassium.

### Iron Deficiency Anemia

DANA C. MITCHELL, JR., M.D., AND O. B. MAYER, M.D.\*

Columbia, S.C. U.S. Army Hospital, Fort Jackson, S.C.

Until proved otherwise, iron deficiency always means blood loss and may well be the only symptom of a serious disease.

Dana C. Mitchell, Jr., M.D., and O. B. Mayer, M.D., point out that, since iron deficiency is a symptom not a disease, treating the state without searching for the etiology is not justified.

centration are all reduced. In general, a deficiency of iron is the basis of all hypochromic microcytic anemias.

Unless blood loss is evidently external, iron deficiency anemia usually results from gastrointestinal hemorrhage. Stools should be examined repeatedly for occult blood by means of the guaiac test, which

#### MEASUREMENT OF SIZE AND HEMOGLOBIN CONTENT OF RED CORPUSCLES

		Normal
Color index	Hemoglobin, % (gra. per 100 cc. x 6.9) Red cell count, millions per cmm. x 20	100
Mean corpuscular volume	Volume packed red cells, cc. per 1,000 cc. Red cell count, millions per cmm.	86-91 cubic microns
Mean corpuscular hemoglobin	Hemoglobin, gm. per 100 cc. Red cell count, millions per cmm.	27-31 micromicrograms
Mean corpuscular hemoglobin concentration	Hemoglobin, gm. per 100 cc. x 100 Volume packed red cells, cc. per 100 cc.	32-36%

Anemia may be elucidated by determining the red blood cell count, the hemoglobin, and the hematocrit. From these may be calculated the blood indexes (see table), which indicate the size of the red cell, the amount of hemoglobin therein, and the degree of hemoglobin saturation of the individual erythrocyte.

Iron deficiency anemia is typically hypochromic microcytic, and the mean corpuscular volume, the mean corpuscular hemoglobin, and the mean corpuscular hemoglobin conis a simple office procedure and, if positive, denotes significant organic bleeding.

When iron deficiency anemia is the consequence of intermittent occult blood loss, chronic duodenal ulcer, hiatus hernia, and ulcerative carcinoma of the gastrointestinal tract are some of the common causes that must be considered.

After maturity is reached, women require sufficient iron supplies to replace blood lost in menses, while men need only comparatively small

 <sup>★</sup> The diagnosis and clinical significance of iron deficiency anemia. J. South Carolina M. A. 65:343-347, 1950.

amounts. Hence, iron deficiency in a man nearly always means blood loss.

Contrary to previous theories, achlorhydria may aggravate iron deficiency but is rarely the cause.

The most economical and effective method of treatment for iron deficiency is ferrous sulfate, 5 gr., three or four times daily. Liver, folic acid, and vitamin B<sub>12</sub> are not necessary for restoration of iron deficiency, nor does molybdenum, copper, or ascor-

bic acid increase the absorption or utilization of iron.

The body does not readily excrete iron; practically none is lost in the urine; the total loss, less than 1 mg. daily, is in the bile and stool. This amount apparently cannot be exceeded except by bleeding. Hence, excessive quantities should not be given. Intravenous administration is rarely justified and may lead to excessive iron accumulation and, possibly, hemosiderosis.

§ PROCAINE AMIDE (Pronestyl) may prevent arrhythmias occurring with upper abdominal or thoracic surgery or with coronary occlusion. The drug depresses blood pressure less than procaine hydrochloride and is hydrolyzed slowly. J. Murray Kinsman, M.D., and associates at the University of Louisville, Ky., find that procaine amide converts ectopic ventricular contractions and paroxysmal ventricular tachycardia to normal sinus rhythm. Nodal tachycardia and ectopic auricular contractions are apparently relieved to some extent by procaine amide.

1. Kentucky M. A. 48:509-511, 1950.

§ FUNNEL CHEST produces typical changes in electrocardiograms. In 8 of 13 cases, William Dressler, M.D., of New York City and Hugo Roesler, M.D., of Philadelphia, found inverted or notched T deflections in multiple precordial leads and also noted the patterns of right ventricular strain. With deep sternal depression and cardiac displacement to the left, features of lead 1 resembled those of dextrocardia. Precordial pain and T inversion in right precordial leads might therefore be wrongly attributed to coronary disease.

Am. Heart J. 40:877-881, 1950.

§ ATYPICAL PNEUMONIA subsides with aureomycin therapy more rapidly than with sulfonamide or penicillin. Effects of the drugs were compared by Emanuel B. Schoenbach, M.D., of Johns Hopkins University, Baltimore, and associates. In 22 cases observed at Sinai Hospital, Baltimore, in 1946 and 1947, the average febrile period with the earlier antibiotics, alone or together, averaged thirteen days from onset. Aureomycin given to 33 patients in 1948 and 1949 shortened duration of fever to about eight days.

New England J. Med. 241 799-806, 1940.

#### Management of Bacterial Pneumonias

HARRISON F. FLIPPIN, M.D., AND HAROLD L. ISRAEL, M.D.\* University of Pennsylvania, Philadelphia

Pand of less training inexpensive and of low toxicity, is the most effective antibiotic against pneumonia caused by pneumococci and other cocci and for most cases of the discase caused by other bacteria.

Aureomycin, chloramphenicol, or terramycin may be considered as the drug to use for infections due to Hemophilus influenzae, Klebsiella pneumoniae, Bacillus tularensis, and Brucella, as well as mixed infection. Moreover, because of the difficulty in differentiating bacterial from viral infections, against which penicillin is ineffective, the tendency is increasing to employ one of these antibiotics as initial treatment for all pneumonia.

Although the pneumococcus remains the most common invader of the lungs, the incidence of classical pneumococcal pneumonia is diminishing, observe Harrison F. Flippin, M.D., and Harold L. Israel, M.D. Meanwhile, pneumonia from gramnegative bacteria, viruses, and rickettsiae is increasing in frequency.

Streptomycin, uniquely valuable for tuberculous pneumonia, is useful for therapy of Friedländer's, tularemic, and influenzal pneumonia and may be an adjuvant to other antibiotics in the treatment of patients with pulmonary brucellosis.

Typhoid pneumonia should preferably be treated by chlorampheni-\* Management of bacterial pneumonias. M. Clin. North America 34:1653-1666, 1950.

col, which is attended by fewer gastrointestinal side effects than aureomycin.

#### DIAGNOSIS

Pneumococcal pneumonia is distinguished from other pulmonary infections by the explosive onset, more massive lung involvement, and relatively rapid resolution of the disease. Roentgenograms are desirable demonstrate the extent and intensity, not the etiology of the involvement. Multiple cavities seen by roentgenology are frequently caused by staphylococcic or Klebsiella pneumonia.

The isolated organism should be tested against the specific therapeutic agents. Blood culture, obtained before therapy is instituted, may identify the causative organism when many bacteria are found in the sputum.

Total and differential white blood cell counts should be made preferably before therapy is started. A high initial white cell count which diminishes rather rapidly after forty-eight hours of treatment is usually of good prognostic import, just as the failure of a low count to rise is generally a poor prognostic sign. Although not diagnostic, the white blood cell count is often helpful in distinguishing bacterial from viral infection.

Therapy should be instituted as

soon as the clinical diagnosis is made and blood and sputum samples have been collected. Antibiotics may be changed when the specific organisms have been determined.

#### THERAPY

Penicillin-Four penicillin preparations are commonly employed:

1 Crystalline benzyl penicillin in aqueous solution, 100,000 units administered intramuscularly, reaches a peak plasma concentration of 3 units fifteen to thirty minutes after administration and maintains a therapeutic level for three hours. In treating bacterial pneumonias, injections should be made every eight to twelve hours. This method of dosage is cheap and rapid acting, has the shortest duration of toxic reactions and the highest peak level with consequent greatest diffusion into serous cavities and abscesses, but is inconvenient for home use since two to three daily injections are required.

2 Procaine penicillin with sodium carboxymethylcellulose in an aqueous suspension (Crystacillin), 300,000 units administered intramuscularly, has a peak level of 1 unit two hours after injection with a therapeutic level lasting twenty-four hours. Doses are given every twelve to twenty-four hours. Most economical for routine hospital use, this preparation requires only one daily injection for most infections but does not yield high plasma levels.

3] Procaine penicillin with aluminum monostearate in oil, 300,000 units administered intramuscularly, attains a peak level of 0.5 units one hour after injection and main-

tains a therapeutic concentration for forty-eight hours, frequently for ninety-six. Injections are given forty-eight hours apart. Although most convenient for home administration, this preparation occasionally produces prolonged toxic and allergic reactions to the penicillin, procaine, or oil.

4] Penicillin, buffered, 200.000 units in oral tablets, yields a peak of 0.4 units thirty minutes after administration, with the therapeutic level lasting three hours. Tablets are taken every three hours. Although easily administered, this preparation is expensive and causes toxic reactions more frequently than the others.

The parenteral route is the most reliable during the first forty-eight hours of therapy for pneumonia, since some oral preparations are not absorbed consistently. The two penicillin schedules which seem most useful in treatment of pneumonia are the aqueous form intramuscularly in 100,000-unit doses every eight to twelve hours, or procaine penicillin in daily injections.

Streptomycin—When treating acute pneumonia for short periods, streptomycin may be used in doses of 0.5 gm. four times daily without fear of toxicity. For tuberculous pneumonia, 1 gm. a day is adequate and is administered in combination with para-aminosalicylic acid, which retards the emergence of streptomycin-resistant strains.

Aureomycin—The gastrointestinal irritation produced by oral use of aureomycin may necessitate lowering the dosage to ineffective levels or discontinuing the drug. Antacid and

aluminum hydroxide preparations, which help to alleviate the gastrointestinal distress, materially lessen the plasma concentrations of aureomycin.

For bacterial pneumonia, the usual dosage of aureomycin is 1 to 2 gm. daily for adults, in divided doses at six-hour intervals. The antibiotic may be administered intravenously in doses of 0.5 gm. every twelve hours. The buffered solution is either injected slowly, directly or added to an infusion of isotonic saline or glucose.

Chloramphenicol—When given to seriously ill patients, an initial loading dose of 60 mg. of chloramphenicol per kilogram of body weight may be given in two or three parts at hourly intervals. Thereafter, 250 mg. is taken every three to four hours for as long as a week after the patient is afebrile. With less severe pneumonia, 25 mg. per kilogram of body weight daily is adequate, in doses of 250 mg. every four to six hours for four or five days.

Terramycin—The dosage of terramycin is 250 mg. every four to six hours.

Sulfonamides—The sulfonamides are largely relegated to pediatric practice, since serious complications, especially periarteritis nodosa and lower nephron nephrosis, may occur when such drugs are given to adults. Used mainly as adjuvants to the antibiotics for desperately ill patients, the sulfonamides are best given in combination, either a mixture of sulfadiazine and sulfamerazine, or a triple mixture of these and sulfamethazine.

#### COMPLICATIONS

Small and moderate sized pleural effusions are not infrequent during therapy and are sterile. Many of these effusions will resorb with continued systemic antibiotics.

Large accumulations of fluid, however, require repeated thoracentesis, and, if persistent, may be treated only by obliteration of the pleural space with resultant expansion of the lung. If the lung has become incarcerated in a fibrinous envelope, either surgical decortication or the fibrinolytic enzyme, streptodornase, should be used. The latter is introduced into the effusion and allowed to exert fibrinolytic activity for twenty-four hours, at which time a thorough aspiration is performed.

Current management of empyema consists of daily aspiration and instillation of 50,000 units of penicillin in addition to high level systemic penicillin therapy. If the lung does not expand, enzymatic decortication with streptodornase should be attempted. When medical therapy is unsuccessful, surgical decortication is indicated.

Most lung abscesses complicating pneumonia can be eradicated by systemic antibiotic therapy employing penicillin or other antibiotics in high dosage. If these methods fail to close the abscess cavity, thoracotomy and pulmonary resection may be done with greater safety and better preservation of pulmonary function than could have been achieved by surgical drainage.

Meningitis, a relatively uncommon complication of pneumonia, may require intrathecal penicillin for cure. If acute bacterial endocarditis complicates the pneumonic process, adequate doses of antibiotics should be administered before extensive valvular damage occurs, and chemotherapy should be continued for at least four weeks. Purulent pericarditis is treated by local instillation of penicillin as well as systemic administration in high dosage. The characteristic electrocardiographic pattern of pneumococcal pericarditis shows extremely high ST segments.

#### Liver Tests in Congestive Heart Failure

LEONARD FELDER, M.D., ALVIN MUND, M.D.,
AND JULIUS G. PARKER, M.D.\*

DEFINITE impairment of liver function occurs with congestive heart failure, but the pattern is not pathognomonic. The exact basis for the change is unknown.

Ten liver function tests were evaluated in a study of 135 cases of protracted congestive heart failure by Leonard Felder, M.D., Alvin Mund, M.D., and Julius G. Parker, M.D., of Montefiore Hospital, Bronx. Two or more different tests were performed for each patient, and half the patients were given at least seven. The bromsulfalein and serum bilirubin determinations were found to be the most sensitive.

No significant differences were noted when the etiology of the heart disease was considered. Duration of heart failure or type of rhythm did not alter the results nor did the nutritional status of the patients produce significant change.

Bromsulfalein clearance was abnormal in 85% of the determinations, while thymol turbidity was found to be so in only 31%.

The alkaline phosphatase was elevated in 46% of all determinations and in 83% of the cases of cardiac cirrhosis autopsied.

Serum bilirubin was increased in 52% of the tests performed; 60% of the patients with cardiac cirrhosis had abnormal values.

With serum cholesterol, a significant difference existed between the rheumatic and nonrheumatic persons; the former had 55% abnormal values and the latter only 28%. Cholesterol ester findings tend to parallel the cholesterol results.

The total plasma proteins were quantitatively abnormal in 29% of determinations; 26% of the patients had less than 3 gm. of albumin per 100 cc. Plasma globulin was elevated in 18%.

Cephalin flocculation was abnormal in 22% of determinations and showed a slight increment for the malnourished patients.

\$ Liver function tests in chronic congestive heart failure. Circulation 2:286-297,

# Hematology in Atomic Bomb Syndrome

GEORGE V. LE ROY, M.D.\*
Northwestern University, Chicago

THE most dependable guide to diagnosis and treatment of human victims of atomic bomb radiation is the changing leukocyte count. In an emergency, additional tests may be limited to determinations of hemoglobin, hematocrit, and protein by the copper sulfate method.

The course and severity of radiation injury generally correspond with the blood picture, particularly with leukopenia. The important symptoms become evident about the time of the lowest white cell count. Japanese physicians believe that recovery is unlikely when leukocytes are fewer than 500 per cubic millimeter.

Onset and extent of a hemorrhagic tendency closely follow thrombocyte depletion, which also parallels leukopenia. Red blood cells vary less consistently and are less useful for diagnosis or prognosis.

In the interests of national defense, permission has been granted to publish the detailed hematologic report of the Joint Commission formed by the U.S. Army and Navy to review atomic bomb effects in Japan. Data of special importance are summarized by George V. LeRoy, M.D., consultant to the U.S. Atomic Energy Commission.

Material was gathered from three sources: early blood examinations of casualties by Japanese physicians, ex
\* Hematology of atomic bomb casualties. Arch. Int. Med. 86:691-710, 1950.

amination of survivors by the Joint Commission, and hematologic surveys in hospitals. Although some work was done under extremely unfavorable conditions, results obtained by the various groups agreed remarkably well.

Subjects were classified in two main groups, [1] those with radiation injury evident from symptoms, and [2] patients who were exposed to comparable amounts of radiation and lived at least twenty days.

Radiation injury is shown by two specific signs, epilation and purpura, and by three important suggestive symptoms: vomiting on the day of bombing; mouth and throat lesions, such as ulceration and gingivitis within thirty-nine days after exposure; and hemorrhagic manifestations other than purpura.

Degree of exposure can also be judged by distance from the point directly under the bomb and by type of shelter. People outdoors or in wooden buildings within 1,100 yd. of the explosion center usually die within a week or two, and 50% of those between 1,100 and 1,650 yd. away die in three to six weeks. At a distance of 2,200 yd., injury is slight.

Serial leukocyte counts of severely injured patients in hospitals agree with those of the same type classified by symptoms or by distance from the bomb. Leukopenia is most pronounced in the third to fifth weeks after exposure and disappears during the eighth to tenth weeks.

In those who die of radiation injury in the first two weeks, however, leukocyte counts usually drop below 1,000 and may be 50 or less.

The number of thrombocytes of seriously affected persons average about 105,000 during the fourth week, when bleeding time is twelve minutes, forty-five seconds and clotting time six and a half minutes. In

fatal cases the level drops to 50,000 or less in the third to sixth weeks.

The lowest hemoglobin values are noted in the sixth to eighth weeks and rising levels after the ninth week. Red cells are enlarged during recovery, probably owing to reticulocytosis.

Among those gravely ill who live at least twenty days, epilation begins about the twentieth day, and purpura on the twenty-fifth. The third to fifth weeks are critical.

#### Rapid Warming and Prolonged Massage for Frostbite

WAYNE G. BRANDSTADT, M.D.º

Ressian investigators, in a symposium on frostbite, advise several measures traditionally banned. Col. Wayne G. Brandstadt, M.C., U.S.A., Washington, D.C., reviews the methods.

Flesh of living men almost never actually freezes. Frostbite results from prolonged low temperature but does not reach beyond ankle and wrist joints, and blood changes are due to venous stasis. The more rapidly a frostbitten extremity is rewarmed, the less necrosis will develop.

Therefore, affected tissue should be rewarmed as quickly as possible by baths and massage. Since overheating is dangerous, temperature of the bath should never exceed 98.6° F. Artificial respiration may be required.

Hyperemia is encouraged by intravenous injection of 10% calcium chloride solution, a total of 15 cc. in 5-cc. doses at intervals of ten minutes. Hot drinks are administered.

Prolonged massage is given in a way that avoids injury. The skin is wiped with alcohol swabs, the body is tightly wrapped in a sterile towel, then vigorously rubbed through the towel with the open palm. If treatment is delayed for several hours, inflammation develops and tissues need rewarming and rest with no massage.

After initial therapy of first-degree frostbite, the skin is rinsed with 5% tannic acid-alcohol solution or painted with 2% solution of brilliant green. During the first three or four days, a sunlamp is used for ten- or fifteen-minute exposures.

# Frostbite, Mil. Surgeon 107:386-388, 1050.



Photos By Courtesy of British Information Services

STRICT ASEPSIS is observed in dressing burns. This view shows the air-conditioned dressing station at Birmingham, England, through a window in the sterilizing room.

# Burns Unit in British Hospital Gets Results


Dissatisfied with results being obtained in the treatment of burns, the Medical Research Council of Great Britain, in 1944, established a Burns Research Unit at the Accident Hospital in Birmingham, England.

After six years some of the answers are still missing. But progress has been made and the unit is continuing its study in an effort to effect speedier and more satisfactory results in the treatment of burns and scalds.

Measurable benefits have already been obtained. Mortality from severe burns has been reduced. Healing time has been shortened. Functional and cosmetic results have been progressively improved. These achievements have been brought about, not by revolutionary discoveries, but by attention to details, refinement of technics, and vigorous application of lessons learned through painstaking research and the analysis of thousands of case records.

To head the original unit with

gists. biochemists, and nurses to evaluate the current methods of treatment and to develop means of obtaining better results.



AIRBORNE INFECTION is prevented by a specially designed dressing room. Air from a roof vent, filtered, warmed, and moistened, is introduced through ceiling vents. An exhaust fan draws stale air under the door and out through the other side of the corridor, thus removing any organisms liberated by the loosening of dressings.

thirty-eight beds and cots at Accident Hospital, the Medical Research Council selected Dr. Leonard Colebrook who had already gained some renown by his pioneering of the use of sulfonamides for the control of puerperal sepsis.

Dr. Colebrook organized a research team of surgeons, bacterioloThe gravest danger to the life of a severely burned patient is the initial shock. This must be counteracted immediately. Transfusions of human plasma effectively maintain the normal circulation of the blood and replace fluids lost from the body. When human plasma is not available, Dextran, a manufactured pro-

duct, may be used. The Burns Unit has found Dextran an excellent substitute.

Apart from the fluid lost during the period of the initial shock from are infected, is prevented by blood transfusions.

When antishock therapy is successful, infection is a big problem. From the first, Dr. Colebrook's team



TRANSFUSIONS are essential to combat shock. Dextran is used as a plasma substitute at the Accident Hospital. Here a doctor is testing the viscosity of a new batch of Dextran with a viscometer suspended in a thermostat bath.

a severe burn or scald, the burn oozes continuously until healed. With a large burn, much of the body's store of protein may be lost. This deficit is replaced by a diet rich in protein foods, such as meat, milk, and eggs. Otherwise the patient loses weight and becomes weakened. Anemia, a common complication of burns, especially if burns

attempted to treat burns under conditions which would eliminate both contact and air-borne infections. Dr. Colebrook, believing that most of the sepsis of burns results from infection in the hospital and can be prevented if the proper conditions and the appropriate routine are established, set up two objectives: [1]

(Continued on page 81)








GRAFTING is not essential with all severe burns but, when necessary, is most effective if done promptly. Formerly a delay of three or four weeks was required to distinguish between burned areas that would heal spontaneously and those that would require grafting. Now, by determining the sensation to a pinprick, decision as to treatment may be made at the first examination. At the left the use of the apparatus for the pinprick test is demonstrated. A sensation-testing needle is used for immediate determination of the depth of skin loss. From the degree of impairment of sensation the physician is able to decide whether grafting is necessary before the slough. The prick can-not be felt if the burn has destroyed the nerve endings. If sensation is gone, grafting is done at

SPARE GRAFTS are not discarded but are stored in a refrigerator to be used later if necessary. The skin remains viable for several weeks. Grafts from each donor are separately labeled and dated and are ready for immediate use if the area of total loss is more extensive than was initially apparent. The bottles containing the donor skin are arranged in racks similar to the one the technician at the right is holding.

Photos on pages 75 to 79 supplied by the British Information Services





GOOD MORALE is an important consideration and may be helped by amenities such as an occasional glass of beer. The man above has a tube pedicle graft which has been raised on his back and transferred to his arm.

to block transmission of pathogens to burns at the time of dressing, and [2] to block transmission of organisms to the burns between one dressing and the next.


By 1948, the first of the objectives


was attained. A special room ventilated by an abundant stream of filtered air was built for the dressing of burns. The dressing station was supplied constantly with filtered bac-

(Continued on page 86)









teria-free air from an air-conditioning plant. An exhaust fan was installed to carry away any dangerous organisms liberated into the air during the removal of soiled dressings. Thus the air in the dressing station is always pure and clean when the next patient arrives. All dressings are done by a trained team using a strict aseptic technic.

Of the 734 patients treated in the dressing station through 1948, less than 1% were infected with the hemolytic streptococcus and Pseudomonas pyocyanea at the time of dressing.

To prevent infection while the patient is in the ward, a layer of penicillin cream is spread on the burn and covered by a thick protective wrapping of gauze and cotton, securely bandaged to form a dressing almost impenetrable by bacteria.

The majority of infections transmitted between one dressing and the next occur with burns difficult or impossible to cover adequately. Another source of infection is the dressing soaked with serous exudate during the first few days after burning. Pathogens conveyed to the outer dressing from the air or bedding are able to grow through the dressing and so infect the burned areas.

In the laboratories attached to the Burns Unit, much research work is being done to devise a dressing that will seal a burn against infection. One such dressing is nylon film which will allow perspiration from the skin to evaporate without permitting the entry of moisture and infection to the burn. The use of cellophane as a barrier to the growing through of

the bacteria has also been suggested.

Because infection has not, thus far, been entirely prevented, chemotherapeutic control of infection is of great importance. Investigations are pursued constantly in the laboratories of the unit to find the antibiotics most effective against the organisms likely to infect burns. Streptococcic infections have proved to be the least difficult to control because they can be quickly eliminated by local application of penicillin or the sulfonamides. Polymyxin is being used with some success against Ps. pyocyanea which sometimes causes harmful infection.

When infection is prevented or eliminated, superficial burns, in which only the partial thickness of the skin is burned through, heal rapidly.

In other cases, prompt surgical attention is essential to good recovery of function. When the complete thickness of the skin has been destroyed, the burn or scald must be grafted if healing is to be quick and deformity prevented. Grafts are taken from unburned areas of skin. Donor areas heal rapidly, leaving no scars.

Grafting is done as soon as the surgeon can distinguish between the partial loss areas, which will heal spontaneously, and the complete loss areas, which will not. Formerly, differentiation of these areas had to wait three to four weeks until the dead skin had sloughed naturally, leaving a raw area behind.

Now, however, Dr. John Bull has found a method which helps in immediately distinguishing one from the other by testing the reaction of the burned skin to a pinprick. In a deep burn the prick of the pin cannot be felt because the nerve endings in the skin are destroyed; in a superficial burn the nerve endings are intact and the pin can be felt. Such knowledge is of the greatest assistance to the surgeons who are enabled in many instances to identify and remove areas of dead skin on the day the burn occurs and to proceed at once with skin grafting. In this way healing is greatly speeded.

At present, the Burns Unit at Birmingham is unique in Britain. Eventually similar projects are expected to be developed in other city health centers. Meanwhile work continues at Birmingham to lay down standards of treatment which other units might follow. Dr. Colebrook retired in 1948, but the investigations he initiated are being continued under his successor, Dr. John R. Squire, Leith Professor of Experimental Pathology, University of Birmingham.

#### Surgical Treatment of Gigantic Hernias

SIDNEY E. ZIFFREN, M.D., AND NATHAN A. WOMACK, M.D.\*

CREATION of an abdominal incisional hernia before definitive repair of gigantic hernias is proposed by Sidney E. Ziffren, M.D., and Nathan Womack, M.D., of the State University of Iowa, Iowa City.

Formation of such a temporary hernia allows the peritoneum to stretch, creating space, and prevents the respiratory difficulty and circulatory embarrassment so often resulting from pressure upon the diaphragm and on the portal or renal venous systems when a large mass of bowel is replaced into the abdominal cavity. After adjustment to pressure, the incisional hernia is closed.

A long transverse incision is made in the upper abdomen through skin, fascia, muscles, and transversalis fascia down to the peritoneum, allowing the peritoneum to bulge into the wound. The skin incision is then closed with a continuous subcuticular suture of No. 0000 steel wire and some interrupted No. 0000 silk. Bowel and omentum of a huge inguinal, umbilical, or diaphragmatic hernia are then easily replaced into the abdominal cavity, and repair of the hernia is done. With inguinal hernias, the testicle on the involved side may be removed and the internal ring closed tightly, with reinforcement of imbricated fascia.

Approximately twelve days later, a nasal tube is passed into the small bowel and the incisional hernia is repaired. The margins of the fascia are easily brought together with No. 000 steel wire. The nasal tube is removed on the second postoperative day.

<sup>\*</sup> An operative approach to the treatment of gigantic hernias. Surg., Gynec. & Obst. 91:709-710, 1950.

### Surgical Management of Indurated Leg

EUGENE L. LOWENBERG, M.D.\*

Norfolk General and De Paul hospitals, Norfolk, Va.

Conservative therapy has little value in cases of chronic indurated cellulitis of the leg, the end result of lymphedema from recurrent thrombophlebitis, lymphangitis, or local trauma.

Eugene L. Lowenberg, M.D., removes widely all the abnormal structures of the involved area, including subcutaneous tissue, fascia, and devitalized or ulcerated skin, and covers the defect with a graft resting on well-vascularized muscle. Any existing venous stasis is corrected simultaneously.

With relatively healthy overlying shin—A longitudinal skin incision is made across the center of the induration from the lower to the upper limits and carried down through the subcutaneous tissue and fascia, exposing the underlying muscle (Fig. 14).

By a line of cleavage just within the skin edge, the entire segment of diseased fat and fascia is excised by blunt and sharp dissection (Fig. 1b). Any incompetent perforating veins encountered are clamped, divided, and ligated. Penrose tissue drains are brought out through stab wounds in the posterior flap. The skin edges are trimmed back to viable tissue and approximated with interrupted sutures of silk (Fig. 1c).

The dressing consists of a thick

roll of gauze from foot to knee and an Ace bandage. The drains are removed in forty-eight hours, the sutures in about ten days. Patients may be ambulatory in about a week.

With unhealthy overlying skin— Any abnormal overlying skin is always removed and skin edges are tasked about the excised area to the muscles beneath.

A split-thickness graft from the thigh is used to cover the defect. The grafted area is dressed with Xeroform gauze, flat gauze moistened in saline solution, and a wide piece of sponge rubber for even pressure. The leg is immobilized for about a week in a foot-to-knee posterior splint. The dressing is changed on the seventh or eighth day, and healing is usually complete by the fourteenth.

With indolent leg ulcer—Preoperative treatment with parenteral penicillin supplemented by penicillin-urethane solution or tyrothricin solution locally is usually necessary for associated chronic ulceration. The ulcer is covered with gauze and the entire area, with surrounding skin, is excised (Fig. 2).

After the operation, the dressing should be kept moist by injecting glycerine, 15%, acetic acid, 0.5%, saline solution every eight hours through catheters enclosed in the

The surgical management of chronic indurated cellulitis of the lower extremity (the indurated leg). Surgery 28:852-850, 1950.

Operation for Indurated Leg  $\boldsymbol{a}$ 

FIGURE 1

b

dressing. Pyocyaneus infection is thus avoided.

With exposed bone and indolent leg ulcer-When unscarred periosteum lies in the ulcer region, the graft is applied directly to the periosteum. Although part of the graft may die, an excellent covering is afforded the bone during the formation of granulation tissue and, later, the dead portion of the graft may be excised and a second graft applied, usually with success.

If bone is diseased, the periosteum sopened sufficiently to remove the

veins are also incompetent, a concomitant ligation of the superficial femoral vein is done, if the venous pressure in the vein is not above go cm. in saline solution. If the pressure is above 30 cm., either the saphenous vein or the superficial




Fig. 2. Indurated leg and indolent ulcer

cortex in layers until normal bone appears. Grafting is then done directly to the bone.

With varicose veins-The saphenous vein is ligated at the femorosaphenous junction when varicose veins occur with chronic indurated cellulitis. After division, the vein is stripped to just below knee level. since the lower branches are resected as the indurated area in the lower leg is removed.

When deep and communicating

femoral vein is ligated, but not both veins.

Postoperatively, edema and venous stasis may be prevented by elevating the foot of the bed about

Anticoagulant therapy is begun four to six hours after surgery. When the patient starts walking, success depends on the prevention of leg edema, so a wide Ace bandage is worn until the circulation becomes competent.

# Hemorrhage from Esophageal Varices

HOWARD K. GRAY, M.D., AND FRANK B. WHITESELL, JR., M.D.\*

Mayo Clinic, Rochester, Minn.


ATTEMPTS to stop bleeding from esophageal varices have been generally unsatisfactory, but a combination of splenectomy, devascularization of the lower part of the esophagus and the cardia of the stomach, and bilateral vagotomy with gastroenterostomy may Azygos succeed.

The usual surgical methods include splenectomy to reduce the volume of blood entering the portal system, removal or injection of existing varices over the lower esophagus and cardia of the stomach, and various shortcircuiting maneuvers to decompress

the portal bed by shifting the circulatory burden to the inferior vena cava or a tributary.

Howard K. Gray, M.D., and Frank
B. Whitesell, Jr., M.D., point out
that emphasis has been unduly placed
on the portal system in treating
esophageal varices to the neglect of

\* Hemorrhage from esophageal varices. Ann. Surg. 152:798-810. 1950.



the azygos system (see illustration).

The portal pressure, spleen, and liver may appear normal, indicating that the primary origin of the varices may be abnormality of the azygos system. Passive congestion or stasis in the azygos veins, lack of valves, poor support afforded by the loose surrounding stroma. and inherent certain structural defects could produce varices despite normal pressure gradients in the portal system. The erosive action of regurgitated gastric acid chyme may precipitate hemorrhage.

In the recom-

mended surgical therapy, devascularization denotes only the stripping of any large varicosed periesophageal veins and the ligation and excision of the arterial trunks ramifying outside the muscularis, the procedure being extended to include a significant portion of the veins over the cardia

Modern Medicine, Feb. 15, 1951

of the stomach. Removal of the extramural vessels does not compromise the viability of the esophagus and abolishes channels which augment stasis and engorgement of dilated submucosal vessels.

Bilateral vagotomy abolishes the

cephalic phase of gastric secretion and reduces gastric acidity and regurgitation of acid chyme into the lower part of the esophagus. Gastroenterostomy compensates for the decreased motility of the stomach following section of the vagi.

§ ODORS FROM COLOSTOMY may be abolished by a mixture of chlorophyll and kaolin inserted in the bowel opening. Julien M. Goodman, M.D., of Ohio State University, Columbus, gave plain gelatin capsules, each containing chlorophyll and 360 mg, of colloidal kaolin, to 18 patients with permanent colostomies. Apparently a single capsule, placed within the colostomy immediately after the morning irrigation and evacuation effectively deodorizes for several hours. The chlorophyll-kaolin mixture produced no undesirable change in character of stool or in bowel habits or other untoward side effects. Patients were able to pursue usual activities without fear of embarrassment.

Surgery 28:550-551, 1950.

§ SURGICAL SCRUB with hexachlorophene is most effective if preceded by a two-minute application of soap and brush. The hands should then be immersed for two minutes in a solution of 0.1% hexachlorophene and 0.5% cetyl alcohol in 70%, by volume, of isopropyl alcohol. The method was thoroughly tested by Irvin H. Blank, Ph.D., and associates of Harvard University and the Massachusetts General Hospital, Boston, under laboratory and actual operating conditions. Rapid degermation is not achieved without the preliminary soap scrub. After repeated scrubbing, the effect of hexachlorophene is accumulative. Cetyl alcohol is emollient.


Surg., Gynec. & Olist. 91:577-584, 1950.

§ BILIARY INFECTION may be treated more effectively by aureomycin than by any other antibiotic. Single intravenous injections of 250 mg, produce levels of 40 to 80 μg, per cubic centimeter in bile, about 6 to 8 times the usual concentrations in serum. Jerry Zaslow, M.D., Maj. Thomas H. Hewlett, M.C., and Ralph Goldsmith, M.D., of the Jewish Hospital, Philadelphia, warn that the drug is fully active only when bile drains freely. But even with obstruction or poor liver function, aureomycin may protect undamaged areas from an ascending process. Antibiotics are adjuncts to and not substitutes for established measures of therapy for biliary infection.

Gastroenterology 16:479-483, 1950.

# Lumbar Sympathectomy


F. M. AL AKL, M.D. Kings County Hospital, New York



KEEP THIS PICTURE IN MIND

#### SURGICAL TECHNIGRAM





1. (Above) Tilt patient 30 degrees with a sandbag under buttocks and lower ribs. Break table.

2. (Left) Incise skin and subcutaneous fat transversely from rectus edge outward 10 cm. to midpoint of line denoting narrowest distance between costal margin and iliocrest. Continue into internal oblique and transversus muscles down to transversalis fascia. Clamp and coagulate bleeders. Incise transversalis fascia at lateral aspect of incision, exposing preperitoncal fat.

3. (Below) Introduce finger into aperture and displace peritoneum medially.




Modern Medicine, Feb. 15, 1951




4. Open resultant fatty fossa with two retractors, and under direct vision, continue reflection of peritoneum medially toward psoas ridge.

5. (Below) With sponge stick reflect peritoneum toward midline and, watching peritoneal fold, roll off psoas muscle and adepoareolar pad between the muscle edge and vena cava or aorta, as the side may be. 6. (Below) Separate transversalis fascia from peritoneum anteriorly. Place wet pad over and retract peritoneum medially. Hold peritoneum with wide Deaver retractor and spread and cut fibrous strands across psoas edge and midline.




Modern Medicine, Feb. 15, 1951






 Reflect fat pad containing perivascular lymph nodes with overlying vena cava toward midline, exposing vertebral bodies and sympathetic chain.


9. (*Yelow*) Lift chain, snip communicating branches. Do not mistake blood vessels for nerve filaments; if in doubt, cut between clamps.




8. Reapply wet pad, readjust retractor to rest on vertebral bodies. Tease fascial covering with nerve hook; isolate sympathetic trunk for 1 or 2 cm.; pass ligature carrier beneath and use ligature for traction.


10. (Below) Doubly clamp any vessels crossing chain, cut between clamps, and electrocoagulate ends.






Modern Medicine, Feb. 15, 1951






crossing the chain are encountered, cut chain and pull it over beneath vein.



12. Lift chain and continue dissection caudad until chain disappears over pelvic brim beneath iliac vessels. Apply silver clip to both extremities of chain for roentgen identification of level, then

cut proximal to the clip and save specimen for histologic examination.



13. Inspect field for bleeding. Remove retractors, straighten table, suture the muscle layers, and close skin with vertical mattress sutures.

Commonly, there are three ganglia on each side of the lumbar vertebrae, but, both in number and location, the lumbar sympathetic ganglia and communicating branches are among the most variable and asymmetric structures of the human anatomy. They may be fused into one long spindle on one side or divided into as many as eight beads on the opposite side. The nerve trunks connecting the ganglia also vary in size from one heavy cord to one or more flimsy filaments, and the communicating branches may be one or several.

For practical purposes, the chain is removed between the point of emergence near the diaphragmatic crura down to the point of disappearance beneath the common iliac vessels. This segment usually comprises the essential L2 and L3 ganglia. Dissecting behind the diaphragmatic crus for Li is unnecessary, and when bilaterally accomplished, may result in paralysis of the ejaculatory mechanism in the male. Search beneath the common lliac vessels for L4 may produce venous oozing, and section of L4, which contains postganglionic axones to the sciatic nerve, may greatly increase sensitization of the denervated arteriorlar walls to circulating adrenalin.

The lumbar sympathetic chain may lie exposed over the vertebral bodies or be obscured by fibrous bands. When invisible, the chain may be located by digital palpation over the vertebra at the vertebropsoas sulcus. In some cases the fibrous bands are a dense fascial structure

which renders the chain impalpable. Only after this fibrous layer is teased with a hook can the chain be exposed and isolated.

On the right side, the vena cava may extend onto the psoas muscle and thus overlay the right sympathetic chain completely. The left chain is always more readily exposed when the fatty mass occupying the sulcus between the psoas muscle and the aorta is reflected medially.

Several muscle-splitting incisions are described, but when the table is broken, the overlapping layers of separated muscle fibers crowd one another instead of falling apart as when the muscles are sectioned transversely.

The transversalis fascia and peritoneum are often inseparably adberent anteriorly and separated laterally by preperitoneal fat. If inadvertently incised or opened during the separation, the peritoneum is promptly sutured. The peritoneum is always reflected toward the midline, otherwise the dissecting finger may stray into the lumbar gutter and reach a dead end at the fatty sulcus between the quadratus lumborum and psoas muscles at the side of the vertebral column.

The electrocoagulating current is convenient for sealing bleeding points deep in the wound and may reduce operating time.

The ureter with the spermatic or ovarian vessels is incorporated in, and retracted with, the parietal peritoneal leaf. These structures are easily demonstrable and should not be confused with the lumbar sympathetic chain.

### Analgesia in First Stage of Labor


ARCHIE A. ABRAMS, M.D.\*

Boston University

Paix may be abolished during the greater part of labor by intradermal infiltration of hypogastric, suprapubic, inguinal, and sacroiliac areas.

Effectiveness is simultaneous with administration. The anesthesia does not change uterine tone or freEarly labor pains are actually viscerosensory reflexes referred to the skin of the hypogastric area near the midline, the skin over the symphysis pubis and inguinal ligaments, and the skin over the sacrum and sacroiliac joints.

Irritation produced by uterine con-



Distribution of wheals on abdomen and back

quency, intensity, and duration of uterine contractions.

Sedation is unnecessary and the affe attendant depressing effects, particularly on the infant, are avoided. Spir Babies delivered after intradermal focular infiltration anesthesia are pink in scolor and usually cry immediately. Segreta the stage labor. New England J. Med. 243:636-640, 1950.

tractions and the stretching of surrounding tissues is transmitted by afferent fibers through the eleventh and twelfth thoracic roots to the spinal cord to create an irritable focus. Afferent impulses from the skin enter the abnormally irritable segment and give rise to painful by intradermal infiltration anesthesia in firstsensations which are referred back to the peripheral structures supplied by the eleventh and twelfth thoracic nerves, chiefly the skin of the lower abdomen and of the sacral area of the back.

To interfere with the viscerosensory reflex thus established, Archie A. Abrams, M.D., uses intradermal infiltration of a 1.5% solution of Metycaine with epinephrine added in a dilution of 1:200,000.

Approximately 30 cc. of the solution is infiltrated into the skin of the abdomen, starting at the midline just above the symphysis pubis. Infiltration continues in a linear fashion, first to the left and then to the right above the inguinal ligaments, out to the anterosuperior iliac spines, and up the midline of the abdomen for a distance of 7.5

cm., extending laterally on each side for 2.5 cm. (Fig. a).

If the patient also has back pain, the skin over the upper sacrum and over the sacroiliac joints is infiltrated intracutaneously with an additional 30 cc. (Fig. b). The drug can be reinjected, if necessary, without untoward or systemic reaction, although the effects of one injection sometimes will last as long as six hours.

Over one-half the patients have complete relief of pain and more than one-third experience moderate relief, but all are conscious of contractions. The infiltration can be started as soon as the patient is in labor and has pain, regardless of stage of cervical dilatation, because the progress of labor is not stopped by the anesthesia.

§ VAGINAL HYSTERFCTOMY produces less fever if a penicillin suppository is placed in the vagina twelve to fourteen hours before surgery. Samuel J. Turner, M.D., of the Chicago Medical School employs 100,000 units of crystalline potassium penicillin G in cocoa butter, in addition to routine preparation. Only 7 of 100 patients so treated had postoperative temperature as high as 101.2° F. for two to four days. With routine prophylaxis and no suppositories, incidence of hyperpyrexia was 37% in 56 cases and 35% in a series of 210.

Am. J. Obst. & Gynec. 60:806-812, 1950.

§ TRICHOMONAS VAGINITIS may be quickly corrected by Tetronyl, a surface-active mixture of two quaternary ammonium compounds in sodium carboxymethylcellulose. Organisms disappeared in one week in 73 of 100 cases observed at the Hospital of the University of Maryland, Baltimore, and in 21 of the others within four weeks. J. Mason Hundley, Jr., M.D., and associates apply the powdered form weekly by insufflation from a single-dose container through an open bivalve speculum. A jelly form is also used twice a day at home.

Am. 1. Obst. & Gyner, 60:843-850, 1950.

### Poliomyelitis after Tonsillectomy

GAYLORD W. ANDERSON, M.D., GENEVIEVE ANDERSON, M.P.H., AUDREY E. SKAAR, AND FRANZISKA SANDLER, M.S.\*\*

University of Minnesota, Minneapolis

The risk of contracting poliomyelitis is increased at least threefold if tonsillectomy is performed during an epidemic of the disease. The chances of bulbar infection are times greater for children who have recently had their tonsils removed.

These conclusions were formulated from a study of the 1946 outbreak of poliomyelitis in Minnesota by Gaylord W. Anderson, M.D., Genevieve Anderson, M.P.H., Audrey E. Skaar, and Franziska Sandler, M.S.

Until the present study, data have been lacking to determine the occurrence of poliomyelitis of any type in recently tonsillectomized children as contrasted with others under identical circumstances otherwise who have not recently had tonsils removed. Rates based on all children infected with poliomyelitis regardless of age and irrespective of the period of the outbreak are meaningless, since the probable incidence of the disease varies greatly at different ages and stages of the epidemic cycle. The only truly valid comparisons are those based on groups of equal age and periods of exposure.

Detailed epidemiologic information was obtained concerning 2,709 persons who had poliomyelitis in Minnesota in 1946. Of these, 19 had had tonsils or adenoids removed during the previous month. Of the 19 patients, 12 or 63.2% had the bulbar type of involvement as contrasted with only 4 of the bulbar variety among 18 patients, 22.2%, who had tonsils removed in the second and third months preceding onset.

During the entire outbreak, only 20% of all patients and 11% of those under 12 had bulbar infections. Of the cases in which onset after operation was within the apparent

AITACK RATES IN CHILDREN Minnesota, June 15- July 31, 1946	Tonsillectomized*	Nontonsillectomized
Population 3 to 7 years old	2,686	240.779**
Cases of polio- myelitis	16	191
Attack ratio	$\frac{1/168}{(0.60\%)}$	1/490 (0.20%)
Bulbar cases	1/221	1/2508
Attack ratio	(0.45°c)	(0.04%)

<sup>\*</sup>Tonsills removed during weeks ending 6/15/46 through 7/31/46; developed poliomyelitis during weeks ending 6/29/46 through 8/10/46.
†Developed poliomyelitis during weeks end-

ing 6/29/46 through 8/10/46.
\*\*Average of two estimates.

the risk of poliomyelitis after tonsillectomy. Ann. Otol., Rhin. & Laryng. 59:602-613, 1950.

usual range of incubation period, seven to twenty days, 80% were bul-

A comparison of the attack rate among all children who had tonsillectomies during the preceding month with that of children of comparable age during the same period of time reveals that the risk of contracting recognizable poliomyelitis after a recent tonsillectomy is 3 times greater than otherwise. The bulbar attack rate of the tonsillectomized group is 11 times that of the group without operation.

The attack rates used for the children with recent tonsillectomies are conservative estimates. For instance, case of poliomyelitis following tonsillectomy was omitted because the incubation period was too short to indicate causal relationship, and 2 fatal bulbar cases were omitted because full details were not obtainable.

Also, an additional 2 cases of tonsil removal in August and October were omitted because so few such operations were done in those months that conclusions could hardly be based on a single case in each month. Had all these cases been included, the rate in the tonsillectomized group would have been higher and the differences in the two groups even greater.

#### Fenestration Operation for Otosclerosis

EDWARD H. CAMPBELL, M.D.\$

RESULTS of the fenestration procedure can be foretold to some extent by the patient's age and degree of hearing loss.

Edward H. Campbell, M.D., of the University of Pennsylvania. Philadelphia, finds that improvement is greatest with preoperative loss of 40 to 55 decibels and at ages of 26 to 45 years. For a successful outcome, the original impairment should be at least 35 decibels but not more than 65, so that a 30-decibel rise will restore serviceable hearing.

The average gain for the slightly deaf is 13 decibels and for the extremely deaf 20. If the deficit exceeds 60 decibels, fenestration will help very little. Occasionally, however, a patient with a loss of 40 or more units is completely rehabilitated.

By actual measurement, the group over 45 years of age improve more often than those under 26, although the average gain is about 16 decibels for both classes. Yet initial deafness is less profound in young people, so that useful hearing is regained more often.

With poor bone conduction at the 2,048 level, average post-operative rise is 21 decibels, and with good function 24.5 decibels.

\* Evaluation of the operative indications and results in the fenestration operation for otosclerosis. Arch. Otolaryng. 52:513-532, 1950.

# Diagnosis of Intracranial Aneurysms

ELDRIDGE CAMPBELL, M.D., AND C. W. BURKLUND, M.D.\*

Albany Medical College, N.Y.

Most intracranial arterial aneurysms develop on or near the anterior half of the circle of Willis.


By reason of close anatomic relationship (see illustration), the extraocular and optic nerves are frequently affected. Therefore, the ophthalmologist is often first consulted.

An intracranial aneurysm may per-

sist without any symptoms throughout life or may rupture. Spontaneous sub-arachnoid hemorrhage is usually caused by rupture of an aneurysm. If the rupture occurs within the cavernous sinus, a fistula to the carotid forms.

The intracranial vascular lesion may enlarge without rupture and produce visual symptoms by pressing on the different cranial nerves. The oculomotor, trochlear, and abducens nerves are often compressed by aneurysms of the internal carotid artery. Paresis of all the extraocular muscles on the involved side can develop. Pressure by the aneurysm on the ophthalmic division of the trigeminal nerve may cause pain behind or above the eye.

Oculomotor nerve paralysis characteristically produces ptosis and dilatation of the pupil with little or no pupillary response to light. In



# Ocular manifestations of aneurysms of the circle of Willis. New York State J. Med. 50:2427-2432, 1950.

addition, the patient will be unable to elevate, depress, or adduct the eye. If the abducens nerve is unaffected, the eye will be externally rotated at rest. Diplopia is noted, with the false image projected inward.

Paralysis of the sixth cranial nerve prevents abduction of the eye, and the consequent diplopic projection is outward.

Trochlear nerve injury alone produces only diplopia. The false image is projected down and inward. Optic nerve involvement may induce uniliteral blindness. Homonymous heminopsia results from an aneurysm pressing on the lateral aspect of the optic chiasm.

Aneurysms of the anterior cerebral aftery or the anterior communicating vessel seldom cause extraocular muscle paralysis. However, various forms of hemianopsia, such as bitemporal hemianopsia, may develop from chiasmal compression. Spontaneous rupture is common with aneurysms of these vessels.

Posterior communicating artery aneurysm is rare. The oculomotor nerve is first affected. Ophthalmic nerve pain may also occur. Aneurysms of the posterior cerebral artery behave similarly.

Diagnosis of intracranial arterial aneurysm is best established by cerebral angiography. Treatment is surgical if the location of the aneurysm will allow excision or trapping by clipping the parent vessel.

Before surgery can be attempted, two matters must be carefully determined: the precise location of the aneurysm, and the presence of an adequate collateral circulation from the contralateral internal carotid artery.

At the time of angiography, compression of the opposite carotid artery in the neck will often reveal the extent of collateral routes by the appearance of Diodrast in the opposite carotid tree. The Matas test for collateral circulation is also employed by Eldridge Campbell, M.D., and C. W. Burklund, M.D. This test consists of digital compression of the carotid artery in the neck for ten to twenty minutes. If signs or symptoms of a unilateral cerebral ischemia appear during compression, collateral circulation is inadequate or absent. The Matas test, however, fails to differentiate between an anterior and posterior communicating collateral circulation.

Depending upon the position of the aneurysm and the state of collateral blood flow, the aneurysm may be excised or clipped. Aneurysms of the carotid artery within the cavernous sinus may be treated by clipping the artery beyond the sinus and ligation of the internal carotid in the neck.

An aneurysm of the intracranial internal carotid between the cavernous sinus and the posterior communicating artery may also be trapped. At present, surgical treatment of an aneurysm at the bifurcation of the carotid artery is too dangerous.

Aneurysms of the anterior cerebral artery proximal to the anterior communicating branch are usually amenable to operation. If the posterior cerebral artery is the site of an aneurysm, the occipital lobe must be removed with the aneurysm.

Recovery of ocular motor func-

In the dictionary

"Joint" sometimes means...

"Joint" sometimes means...





On gamma globulin labels

(human)
sometimes means
placental blood

(human) - CUTTER

<u>always</u> means

100% venous blood



#### 100% Venous Blood Means:

- MINIMAL REACTIONS Cutter fractionation process produces a highly concentrated gamma globulin with resultant small dosage, plus crystal clear purity to hold reactions to a minimum.
- CRYSTAL CLARITY-Look at the highly purified homologous protein in the Cutter vial. See the crystal clear, homolysis-free gamma globulin.
- STANDARD POTENCY—Each lot of Cutter globulin is made from human venous blood pooled from 4500 male and female donors to assure consistent anti-measles activity - providing a constant gamma globulin content of 160 mgm. per cc.
- As a product of choice for measles modification or prevention,

insist on

CUTTER Immune Serum Globulir

-the only gamma globulin -available from your pharmacist fractionated entirely from human senous blood

CUTTER LABORATORIES . BERKELEY, CALIFORNIA

Producers of concentrated 2.5 cc. Hypertussis - specific for passive immunity against whooping cough.

tion after excision or trapping of an aneurysm is variable. The trochlear and abducens nerve paresis frequently disappears. Oculomotor nerve damage is less often reversible. Some deficit often persists; misdirection of fibers may also occur. The patient frequently cannot perform third nerve functions, except adduction, with the involved eye.

#### Eczema from Wooden-handled Objects

J. B. HOWELL, M.D., AND D. SHELTON BLAIR, M.D.<sup>©</sup>

RECCURENT eczema of the hands and fingers is occasionally caused by allergic reaction from using or making wooden objects. The only two woods known to produce this vesicular eczema are cocobolo and rosewood, both of which are used for cutlery handles and other utility articles.

Some of the common objects made of these woods are the handles of kitchen utensils, brushes, umbrellas, and small tools, steering wheels, policemen's clubs, canes, billiard cues, rosary beads, buttons, and knobs.

The irritating principle is apparently contained in an oil from the wood. The oil is soluble in alkali but is precipitated by acid. Persons whose perspiration has an alkaline reaction are susceptible, since alkali dissolves the irritating agent and permits entrance into the pores of the skin.

The appearance of cocobolo wood dermatitis of the hands is a protean eczematous eruption, and only by considering all the etiologic possibilities of hand eruptions can the problem be solved. Patch testing of a sensitized person with cocobolo wood shavings may provoke a severe local reaction at the test site and exacerbation or activation of original areas of involvement. Therefore, an ether extract and corn oil dilution of the oleoresin is best for contact testing. Shavings from suspected objects in use for some time seem safe to use in the test because some of the oil has been lost.

J. B. Howell, M.D., and D. Shelton Blair, M.D., of Dallas have recently observed 3 housewives with severe recurrent vesicular eczema of the hands. Results of patch tests for the common causative agents were negative, and the condition resisted all treatment. In every case, patch tests with cocobolo wood shavings gave positive reactions and, after the offending articles were removed from the household, the eczema cleared and did not recur.

Φ Eczema of the hands from wooden-handled objects. Arch. Dermat. & Syph.
62:400-404, 1950.

# convenient and most economical potent androgenic therapy

mucorettes,\* hard compressed disks of methyltestosterone, permit you to maintain patients requiring androgenic therapy "on comparatively small dosage schedules." Placed in the labial pocket formed by the upper lip and gum above incisors, the methyltestosterone is slowly absorbed through the oral mucosa directly into the blood stream.

Methyltestosterone, so administered, is at least twice as effective as that ingested <sup>2,3,4</sup>—on a weight basis—and may in some cases approach the efficacy of testosterone propionate employed parenterally.<sup>4</sup>

Finkler, R. S.: J. Clin. Endocrinol. 7:293, 1947. <sup>\*</sup>Lisser, H.: Northwest Med. 49:949, 1947. <sup>\*</sup>Tyler, E. T.: J.A.M.A. 139:9, 1949. <sup>\*</sup>Escamilla, R. F.: Am. Pract. 3:425, 1949.



# maximum patient economy

For full efficiency, mucorettes should be placed between the upper lip and gum above incisors.

White .

#### mucorettes

Supplied: In disks containing 10 mg. methyltestosterone, scored; in bottles of 30, 100 and 500. rare-galen division of

white laboratories, inc.

pharmacoutical manufacturers newark 7, n. j.

# Intracapsular Hip Fractures

ROBERT T. MC ELVENNY, M.D.\*

Northwestern University, Chicago

I NITIAL loss of blood supply to a bone fragment at time of injury neither prevents nor produces fibrous or delayed union.

Primary avascular necrosis is caused by direct damage to the medullary components of the bone from disease, foreign material, heat, abnormal external or internal pressures or by continued interference with the requirements of bone growth or healing.

Robert T. McElvenny, M.D., believes that conditions such as traumatic dislocation of the hip or injury of the medullary structures of the femoral head compromise the vascular supply. The consequent damage is greater than that caused by disturbance of the extrinsic circulation. The initiating factor may be the crushing or pounding force of the accident, surgical trauma, or the effects of therapy. Delayed union of a fracture is the most common cause of avascular necrosis.


When roentgenograms disclose delayed union, healing may be stimulated by bone graft pegging or osteotomy before avascular necrosis occurs. Continued disunion of a talar head fragment, carpal scaphoid fragment, or capital epiphysis leads ultimately to avascular necrosis, with bone disintegration and poor function. A good bony union of slipped upper femoral epiphysis is induced by fixation with bone pegs more quickly than with metal. When the head is badly displaced, osteotomy is required to restore good position.

In both delayed union and primary avascular necrosis, the first changes noted on roentgenograms are in the femoral head on the acetabular side of the former epiphyseal plate. Evidence of bone production and proliferation appears on the upper proximal portion of the neck when the head substance has diminished sufficiently to permit injury of the neck by the acetabular components.

To diagnose primary avascular necrosis, immediate bony union between the femoral head and neck must be demonstrable and no fracture line visible.

With delayed union, no bony continuity exists between the fragments. Repeated small fractures caused by slight motion of the head and neck produce intermittent attempts at callus repair and continued proliferation of the callus mass, easily detectable on serial roentgenograms (see table). Plates are made every four weeks for the first six months, every six weeks for the next three months, then every three months for two years.

\* An interpretation of certain factors affecting treatment of intracapsular hip fractures. Quart. Bull. Northwestern Univ. M. School 24:267-279, 1950.



#### controlled maintenance...

Through precise control of contractile force and rhythm, Digitaline Nativelle provides positive maintenance of the decompensated heart—maintains the maximum efficiency obtainable. Absorbed completely, it dissipates at a uniformly predictable rate—maintains full digitalis effect between doses with virtually no local side effects. For the comfort and protection of your patients—for your own assurance—specify Digitaline Nativelle in full—on your prescription.

### digitaline nativelle

Chief active principle\* of digitalis purpurea (digitaxin),
\*not an adventitious mixture of glycosides.



For dosage instructions consult Physicians' Desk Reference.

Send for brochure, "Modern Digitalis Therapy,"

Varick Pharmacal Co., Inc., (Division of E. Fougera) 75 Varick St., New York.

#### ORTHOPEDICS

Changes secondary to delayed union include slight condensation about the fracture line, fibrosis of medullary components, cystic changes in the head, sclerosis of the head with calcium precipitation in the solely by functional or cosmetic considerations. For ideal results, reduction should be with the proximal fragment in valgus position, but for rapid bony union, satisfactory reduction is the main concern. Inadequate-

#### ROENTGENOGRAPHIC DIAGNOSTIC TABLE

Primary Avascular Necrosis	Delayed Union			
Head and neck are intact.	Head and neck not united by bone.			
No proliferation is seen about fracture site.	Proliferation of callus at fracture site- usually most marked on superior por- tion of neck just at fracture site. Thi is usually first visible eight to twent weeks postpinning. It may be eviden following a few weeks of full weigh bearing.			
Increased density of head is not marked.	Increased density of head eventually marked.			
Loss of bony substance occurs on head side of old epiphyseal plate line.	Small circular areas of cystic change throughout head eventually appear.			
First loss of bony substance of head is usually seen at upper outer quadrant, centrally if full effect of trauma is delivered there (central dislocation).	Loss of bone substance as gouged-out area in upper outer quadrant of head is not common and, if it appears, is a late sequela.			
Head does not shift or flow over and under inferior medial portion of neck because bony union exists at fracture site.	Head tends early to shift slightly and later flow over and under inferior medial portion of neck. This is not a refracture. The head and neck have never united by bony union.			
Bony union at fracture site is established or continuity between head and neck has never been lost.	Union between head and neck may eventually occur. Head tends to flatten. Changes in head may be obvious at this time; if not, will begin to appear within two years of fracture treatment.			

structural space, and, possibly, disintegration of the head. After a period of months, some cases of delayed union may achieve bony union but, by this time, the head has become necrotic.

Varus and valgus positions are in no way concerned with bony union; the choice of position is governed ly reduced varus hips result in more cases of nonunion than of delayed union. Inadequately reduced valgus hips, except subcapital fractures, give more delayed unions than nonunions because of the favorable thrusts at the fracture site in that position.


Although the degree of angle has



GE MAXICON meets the medical profession's long-felt need for x-ray equipment developed to grow with an expanding practice, providing just the x-ray facility required...unit by unit as needed!



From this ..



to this ... in a few easy steps!

Bucky table to motor-driven combination unit! Comprised of a number of components that can be assembled in various combinations — the Maxicon series have a wealth of utility wherever diagnostic x-ray is employed. The Maxicon covers the range of diagnostic x-ray apparatus from the horizontal x-ray table to the 200-ma, two-tube, motor-driven combination unit.

Check the remarkable flexibility of the Maxicon. Discover what it can do for you. Ask your GE representative for unique booklet demonstration of the Maxicon series, or write General Electric X-Ray Corporation, Dept G-2 Milwaukee 14, Wisconsin.

> GENERAL S ELECTRIC X-RAY CORPORATION

no prognostic value in predicting union, the more oblique the angle, the greater the amount of traction needed and the more difficult to get adequate reduction.

Locked fragments at time of reduction usually produce immediate bony union, while unlocked pieces fail to heal. When reduced, the fragments automatically lock if placed so that the cortex of the femoral head is above and inside that of the neck fragment. Shearing and angulating forces are so converted as to increase the impacting and locking tendencies. This position-the third position-can be defined on roentgenograms in the anteroposterior view with the limb in complete internal rotation, neutral lateral position, and with traction maintained. Traction is applied only to impacted fractures showing inadequate reduction.

The following two types of femoral neck fractures are difficult to reduce and usually result in nonunion or delayed union:

i] A true subcapital fracture with no neck on the head fragment. Extreme difficulty is encountered in attempting to force the femoral neck under and inside the most medial portion of the head to ensure firm locking. Unless this position is attained and maintained after fixation, some form of open reduction is indicated as the initial treatment of choice. Osteotomy may be done or an attempt be made to shape the superior portion of the neck so as to leave superiorly a wide, broad surface for the head to rest on. The procedure is followed by fixation.

2] A fracture in which an abnormally long medial fragment remains on the head. An excellent result may be obtained if the beak fragment is placed lateral to the inner portion of the neck fragment, gouging into the cancellous bone of the neck and wedging firmly lateral to the medial V-shaped cortical defect of the femoral neck. If adequate reduction cannot be maintained, an immediate osteotomy should be done or open operation performed in which the beak is fashioned to invaginate deeply into the cancellous substance of the neck fragment.

§ PHENYLACETYLUREA, Phenurone, is of value in the control of psychomotor epilepsy. Of 24 patients with psychomotor attacks, 18 were improved by Phenurone, report Samuel C. Little, M.D., and R. R. McBryde of the Medical College of Alabama, Birmingham. The daily dose is 2 gm. or less. Slight improvement in grand mal seizures occurs. Little benefit is noted for patients with petit mal attacks. Maximum improvement in seizures is reached during the first two months of therapy. Leukopenia occasionally develops and repeated white blood cell counts must be made during treatment. Toxic symptoms, involving chiefly the central nervous system or digestive tract, require discontinuance of the drug in about one-sixth of cases.

Am. J. M. Sc. 219:494-499, 1950.

# The elastic stocking that "sugarcoats" the prescription

#### One reason more women wear Bauer & Black ELASTIC STOCKINGS is that they are so sheer and inconspicuous

The doctor who considers a woman's pride of appearance when he prescribes an elastic stocking does her—and himself—a service. Because the sheer, unobtrusive Bauer & Black fashioned Elastic Stockings actually flatter the leg, your patient does not object to wearing them.

Bauer & Black Elastic Stockings provide the firm support required for surface varicose veins. Smooth, uniform pressure is insured by their two-way stretch.

#### BAUER & BLACK

# Elastic stockings

Other famous Bauer & Black Elastic Supports: TENSOR\* Elastic Bandage, BRACER\* Supporter Belt, Abdominal Belts, Suspensories, Anklets, Knee Caps, Athletic Supporters. BAUER & BLACK, DIVISION OF THE KENDALL COM-PANY, 2500 S. DEARBORN ST., CHICAGO 16, ILLINOIS



\*Reg. U.S. Pat. Off.

#### Neuropathy in Infectious Mononucleosis

THEODORE C. BERNSTEIN, M.D., AND HAROLD G. WOLFF, M.D.\*

Veterans Administration Hospital, Bronx Cornell University, New York City

THE central nervous system may be affected in infectious mononucleosis. Though the complication is relatively infrequent, some idiopathic acute disorders of the nervous system may be associated with infectious mononucleosis.

The symptoms depend on the site of greatest involvement. Manifestations may appear as serous meningitis, encephalitis, meningoencephalitis, polyneuronitis, or peripheral neuropathy.

The nervous system is implicated in less than 1% of cases of infectious mononucleosis and, as is the case with mumps, neurologic complications are considerably more frequent in males than in females, reports Theodore C. Bernstein, M.D., and Harold G. Wolff, M.D., who review the literature and describe a recent additional case of meningoencephalitis.

Diagnosis of infectious mononucleosis is made on the basis of a heterophil antibody titer above 1:64, the appearance of monocytic blood cells, and cervical or generalized lymphadenopathy.

The common signs and symptoms of nervous system involvement are headache, disturbances of consciousness, nuchal rigidity, convulsions, ataxia, hemiparesis, cranial nerve palsies, papilledema, and motor and

sensory disturbances. First indications usually appear from one to three weeks after onset of the discase, but may be the initial indication of the infection.

The neurologic manifestations are similar to those observed in diseases caused by other acute infectious agents. Therefore, physicians should consider the possibility of infectious mononucleosis in the differential diagnosis of lymphocytic meningitis, encephalitis, and acute polyneuronitis, and heterophil antibodies should be sought in all acute disturbances of the nervous system.

At the present time, infectious mononucleosis is believed to be of virus origin, although this concept is not confirmed. Death occurs infrequently with the disease, and postmortem examinations are seldom done. In the few autopsies that have been performed, alterations in the nervous system were observed. These included meningeal infiltration with mononuclear cells, perivascular hemorrhages, perivascular cuffings of mononuclear cells, alterations of neurons in the brain and spinal cord, and congestion, edema. and cellular infiltration of peripheral nerves and roots.

Spinal fluid changes consist of increase in lymphocytes or protein or both. Abnormalities in the spinal

Involvement of the nervous system in infectious mononucleosis. Ann. Int. Med. 33:1120-1138.



# a most significant advance

# TROMEXAN

ethyl acetate

### new, safer, oral anticoagulant

Throughout the exhaustive studies on TROMEXAN, involving many hundreds of cases, this new anticoagulant has proved singularly free from the dangers of hemorrhagic complication. Other advantageous clinical features of TROMEXAN are:

- 1 more rapid therapeutic response
  (therapeutic prothrombin level in 18-24 hours);
- 2 smooth, even maintenance of prothrombin level within therapeutic limits;
- 3 more rapid return to normal
  (24-48 hours) after cessation of administration.

In medical and surgical practice . . . as a prophylactic as well as a therapeutic agent . . . TROMEXAN extends the scope of anticoagulant treatment by reducing its hazards.

#### Detailed Brochure Sent on Request.

TROMEXAN (brand of ethyl biscoumacetate): available as uncoated scored tablets, 300 mg., bottles of 50 and 250.



GEIGY COMPANY, INC.

Pharmaceutical Division, 89-91 Barclay St., New York 8, N. Y.

fluid may exist without any evidence of nervous system disturbance.

Although infectious mononucleosis is predominantly a disease of children, the nervous system involvement is more common in patients 20 to 30 years old than in younger persons.

Recovery from neurologic abnormalities is complete for 85% of patients. Despite complete symptomatic recovery, spinal fluid changes persist for long periods in many cases. Muscular weakness, wasting, and slight motor aphasia have been reported as permanent sequelae.

#### Serum Cholesterol and Age

ANCEL KEYS, PH.D., AND ASSOCIATES\*

With healthy men, the total serum cholesterol concentration rises yearly until about the age of 55 and thereafter usually decreases.

At all ages above 35 years, values for 1% of males with no discoverable illness exceed 300 mg. per 100 cc.

Serum cholesterol levels were measured at the University of Minnesota, Minneapolis, by Ancel Keys, Ph.D., Olaf Mickelsen, Erma v. O. Miller, E. Russell Hayes, M.D., and Ramona L. Todd, M.D., in two independent research projects.

About 5,000 tests were done of 1,492 men from 17 to 78 years old and of 564 women from 17 to 30 by the Bloor extract Liebermann-Burchard method or the Schoenheimer-Sperry method.

Most subjects were in an upper economic group engaged in business, professional, and scholastic pursuits. Diets were varied and rather luxurious. The influence of basal or total energy metabolism, relative obesity, and food was not considered, however.

Between 17 and 30 years, cholesterol levels of men and women are about the same and increase 2.2 mg. yearly. The average at 22 years is 178.7 for males and 177.8 for females. Men add about 2.3 mg. annually from 17 through 45.

The high male value at 55 years is approximately 255.7 mg., and the mean fall from 60 to 68 is 2.9 mg. yearly. Between ages of 45 and 70 years, 1% of men have levels over 320 mg., and from 45 to 60 years, 5% are over 300 mg.

Although the persons investigated appeared well by careful physical examination, the data represent average, not necessarily the most desirable rates. The steady decline after middle life may indicate demise of those with high values rather than actual decrease. Figures correspond with normal values recently observed in Denmark.

<sup>2</sup> The concentration of cholesterol in the blood serum of normal man and its relation to age, J. Clin. Investigation 20:1347-1353, 1050.

Ammivin is pure khellin. This new weapon against anglna pectoris and coronary insufficiency offers these advantages:

Ammivin is approximately five times as potent a vasodilator as aminophylline.

Ammivin is selective.

Dilates the coronary arteries without appreciable effect on peripheral circulation—does not alter blood pressure •

Each Ammivin tablet contains 20 mg. of khellin to allow flexibility of dosage •

Ammivin is enteric coated .

Ammirin is the preferred dosage form of khellin •

How Supplied: Bottles of 40 and 100 tablets.

Comprehensive Brochure upon request.

## Ammivin

(pure khellin)

a potent coronary vasodilator



The National Drug Company,
Philadelphia 44, Pa.
More than Half a Century
of Service to the Medical Profession

§ SPONGE IN THE ABDOMEN may be recognized in roentgenograms by a small localized collection of air bubbles trapped in the mesh. The pattern is more uniform than that of intestinal gas and is not changed by shift of position or from day to day. A fatal case was diagnosed by M. Slater, M.D., of New York City after development of adhesions, pyloric obstruction, and subphrenic abscess. However, a similar roentgen appearance led to early identification in another instance, in time to save the patient's life.

Am. 1. Roentgenol. 64:781-784, 1950.

#### Localization of Intrathoracic Lesions

BENJAMIN FELSON, M.D., AND HENRY FELSON, M.D.\*

DISTORTION of the usual cardiac silhouette of the posteroanterior roentgenogram is a reliable indication of the location of intrathoracic lesions that are in anatomic contact with the heart or aorta.

Benjamin Felson, M.D., and Henry Felson, M.D., of the University of Cincinnati have determined the accuracy of this new sign in more than 84 cases. Diseases included acute or chronic pneumonia, tuberculosis, atelectasis, encapsulated pleural effusion, and tumor.

The following rules were formulated:

➤ A shadow concealing all or part of the heart border is anterior and, therefore, involves the middle lobe, lingula, anterior mediastinum, lower end of the major interlobar fissure, or anterior part of the pleural cavity.

▶ Density that overlaps but does not hide the heart outline is posterior and thus lies in a lower lobe, the posterior mediastinum,

or the posterior part of the pleural cavity.

▶ Opacity obliterating the right edge of the ascending aorta is in the anterior segment of the right upper lobe, middle lobe, right anterior mediastinum, or anterior portion of the right pleural cavity.

▶ Density which overlies but does not obscure the right border of the ascending aorta is in the upper segment of the right lower lobe, posterior segment of the right upper lobe, posterior mediastinum, or rear of the pleural cavity.

▶ Elimination of the left border of the aortic knob indicates involvement of the apical posterior segment of the left upper lobe

or adjacent mediastinum.

▶ A transparent defect over the aortic knob is either forward in the left upper lobe, high in the left lower lobe, or to the front or rear of the upper mediastinum or pleural cavity.

\$ Localization of intrathoracic lesions by means of the postero-anterior roentgenogram. The silhouette sign. Radiology  $55;363\cdot374,\ 1950.$ 

More physicians in more countries are now prescribing Numotizine because...

... Comforting relief from congestion, pain and swelling in inflammatory conditions is basic.

... Numotizine Cataplasm is externally applied, and is effective over many hours.

... Specific antibiotic and chemotherapeutic therapy is enhanced by the palliative effectiveness of Numotizine.

We shall be pleased to send a clinical trial supply on request.

**Ethically Presented** 

For clinical sample, just write "Numotizine" on your card, letterhead or Rx blank, and mail to us.

NUMOTIZINE, INC.

900 N. Franklin Street

Chicago 10, Illinois

#### Polyps of the Colon

ROBERT TURELL, M.D., AND ROBERT S. WILKINSON, M.D.\*

Beth Israel and Harlem hospitals, New York City

ALL adenomas of the bowel are potentially malignant and so should be removed in toto. Sigmoidoscopic and roentgenographic studies should be made periodically thereafter to detect new or recurrent lesions at the earliest possible moment.

Colonic adenoma appears in 1 of 5 people who reach the age of 60 and in approximately 2% of asymptomatic individuals under the age of 45. Robert Turell, M.D., and Robert S. Wilkinson, M.D., found adenomas in over 7% of 386 patients over 50 years of age who had intestinal symptoms. Of 31 lesions, 5 were malignant and cancer developed after three years in 2 patients who refused treatment.

Polyps of the large bowel occur most frequently in the sigmoid, next often in the ascending colon or rectum, and least often in the cecum and transverse and descending colon.

The first indication of colonic polyps is usually bleeding; next most frequent is protrusion of the polyp through the anus. Sigmoidal adenomas occasionally cause lower abdominal pain.

A comprehensive diagnostic survey includes a bidigital rectal examination, sigmoidoscopy for 25 cm. or farther, biopsy, and barium and air double-contrast roentgenography. The elect is used to Since malignant transformation may adenomas, es Adenomas of the colon and rectum. Surgery 28:651-661, 1950.


occur anywhere in the polyp, an adequate biopsy to detect infiltration or invasion should afford specimens of tissue from the center, the periphery, and the surrounding mucosa, including basal submucosal structures.

A solitary rectal polyp is frequently accompanied by adenomas elsewhere, for which search should always be made.

Pedunculated and sessile benign adenomas and growths with long and narrow pedicles containing noninvasive malignant foci may be removed by surgical diathermy or local surgical excision with a cold or electric scalpel. Except for large polyps or lesions above the peritoneal reflection, most extirpations may be accomplished in the office.

Mucosal excrescences and small sessile or pedunculated adenomas are removed by cold angulated biopsy forceps followed by thorough desiccation of the mucosal base. Even when noninvasive malignant cells are found in the body of the growth, large pedunculated polyps and some medium-sized sessile lesions, especially if located on the posterior aspect of the rectum, may be amputated at the mucosal base by high-frequency snare.

The electric double-loop resector is used to remove large sessile adenomas, especially those below the



a picture of

# Death on Bacteria

No matter how minor the surgery you may perform in your office, you face a grave responsibility. You never can tell when the blood stream of a patient carries spore-bearing bacteria. Consequently there is always the danger of cross-infection when an instrument touches any incision. That calls for the safety of hospital sterilization in your office . . . certain destruction of the toughest bacteria. For such safety, the Pelton HP Autoclave is the perfect answer. It is compact, fully automatic. See it at your dealer's or write for literature.

# **PELTON**

THE PELTON & CRANE CO., DETROIT 2, MICHIGAN

peritoneal reflection, and tumors in inaccessible areas. Invasive polyps are true carcinomas and require radical resection.

Lesions above the reach of the sigmoidoscope are extirpated by transabdominal colotomy when the stalk or pedicle is narrow and long. If the polyp cannot be felt during surgery, transillumination of the gut in a darkened room with a specially constructed, sterile, cold light may be helpful. This failing, the colon may be incised on the antimesenteric side and examined with a sigmoidoscope.

Colotomy through which wide resection of the mucosal base and the submucosal structures may be done is used for the removal of shortpedicled adenomas and small or medium sized sessile lesions without palpable regional lymph nodes. For large sessile polyps, segmental resection is usually required, including end-to-end anastomosis with or without colostomy or ecostomy.

Papillary adenomas are rare, occur usually in elderly patients and in the rectum or rectosigmoid, and feel extremely soft and spongy by digital examination. These lesions recur after treatment and are more often malignant than other colonic polyps. Treatment consists of removal in toto by local excision or electrothermy.

With familial polyposis, malignant changes are nearly always found during the early decades of life. Therapeutic procedure consists of primary fulguration of the adenomas of the lower rectum, anastomosis of the ileum to the lower portion of the rectum, 8 cm. from the anus, for easy subsequent endoscopy, and colectomy in two stages. Total colectomy and proctectomy are indicated for complicating cancer of the rectum or when the adenomas are so numerous as to preclude successful removal proctoscopically.

Isolated adenomas associated with ulcerative or amebic colitis are managed as though unrelated to the inflammatory process, but multiple adenomas complicating chronic ulcerative colitis are best treated by total colectomy because of possible malignant degeneration.

§ PERTUSSIS IN INFANCY may seriously damage the brain. Babies admitted to the Children's Hospital, Boston, for whooping cough at ages between three weeks and two years were examined years later by Randolph K. Byers, M.D., of Harvard University, and Nicholas D. Rizzo, M.D., of Peter Bent Brigham Hospital, Boston. Permanent effects were obvious in 6 of 35 cases not known to have had cerebral defects prior to the pertussis. Findings included feeble-mindedness in 2, mental irregularities implying diffuse cortical injury in 3, and emotional immaturity in 1. In 3 additional cases, transient retardation occurred. Since pertussis vaccine seems prophylactically efficacious, the risks of its use in infancy are probably outweighed by those of the disease.

New England 1, Med. 242:887-891, 1950.

# 1 ST IN THE FIELD! TAR PLUS ANTIHISTAMINE



In many pruritic and allergic skin disorders, the torment of itching is a primary consideration and its alleviation an immediate need. HISTAR'S antihistaminic provides a potent local anesthetic action bringing quick relief of the itching, burning and swelling attending many of these cutaneous disorders.

However, the complete cycle of therapy, as displayed by HISTAR, is fulfilled by the incorporation of tar, universally recognized for its effectiveness in such disorders.

Clinical tests have shown HISTAR to be 72% effective in the improvement, amelioration or disappearance of lesions. In a special study of 54 cases conducted for possible side effects, no evidence of systemic or local

HISTAR—a product of
THE TARBONIS COMPANY
4300 Euclid Avenue, Cleveland 3, Ohio

toxicity was found in the use of HISTAR.

The physiologic-synergistic action of pyrilamine maleate 2% (formerly called pyranisamine maleate) and special extract of coatra 5% assures the patient both comfort and treatment, in one effective ointment.

For prescriptions—all pharmacies stock 2 oz. jars; for dispensing purposes, 1 lb. jars available through your surgical supply dealer.

THE TARBONIS C	OMPANY	Dept. MM
4300 Euclid Avenue	, Cleveland 3	, Ohio
Please send literat HISTAR.	ture and clin	ical sample of
Name		M.D
Address		
Cin	Zone	State

#### Bone Marrow of Infants and Children

KURT GLASER, M.D.

Hadassah Medical Organization, Jerusalem

LOUIS R. LIMARZI, M.D., AND HENRY G. PONCHER, M.D.\*

University of Illinois, Chicago

TELLULAR composition of infants' I bone marrow fluctuates widely during the first year of life, particularly in the first month.

Throughout the rest of childhood and adolescence, cell counts are approximately the same as for adults.

Single tests are unreliable, however, because several aspirations from the same person or even different areas on the same slide may vary greatly. A deviation calls for repeated counts and further investigation.

Much more important than a figure for any type of cell are the relative proportions of various elements and the total morphologic pattern.

Values were established for healthy subjects by Kurt Glaser, M.D., Louis R. Limarzi, M.D., and Henry G. Ponther, M.D. Sternal punctures were done of 42 babies less than a month old, 17 between 1 and 12 months. and 92 children and young people from 2 to 19 years inclusive.

The sternum is pierced in or near the midline of the second or third interspace with a 16-gauge needle. A 1-cc. specimen of marrow fluid is withdrawn, less from small babies. and thoroughly mixed with a little powdered heparin. Centrifugation for five minutes at 2,000 rpm separates the sample into layers.

The myeloid-erythroid stratum is narrow and varies in width during the first year, then becomes more abundant and stable. Fat is seen in traces or not at all below the age of 1 year but is generally measurable thereafter. The layer of mature erythrocytes tends to decrease with advancing age.

Fat and most of the plasma are removed, and marrow smears are prepared with Wright stain. Either 400 or 500 nucleated cells are then counted.

Some errors and misinterpretations are unavoidable with the most careful technic. Any method of aspiration involves dilution by peripheral blood, especially if the bone cavity is very small and the needle point near blood sinuses. Cell counts also differ because of uneven distribution and relatively small numbers.

Erythroid forms are relatively frequent during the first two days of life, diminish until the fifteenth day. and increase to an average of 23.12% by the end of the first month. The level then remains stable except for a temporary fall during the third to fifth months.

The myeloid series also varies con-# Cellular composition of the bone marrow in normal infants and children. Pediatrics 6:789-

824. 1950.



## Poly-Vi-Sol Tri-Vi-Sol Ce-Vi-Sol

for their varying vitamin needs..

	VITAMIN A	VITAMIN D	ASCORBIC ACID	THIAMINE	RIBOFLAVIN	NIACINAMIDE
POLY-VI-SOL each 0.6 cc. supplies:	5000 units	1000 units	50 mg.	1 mg.	0.8 mg.	5 mg.
TRI-VI-SOL each 0.6 cc. supplies:	5000 units	1000 units	50 mg.			
CE-VI-SOL each 0.5 cc. supplies:			50 mg.			

water-soluble liquid citamin preparations...

pleasant tasting, convenient, economical. Available in 15 and 50 cc. bottles with calibrated droppers.





MEAD JOHNSON & CO. EVANSVILLE 21, IND., U. S. A. stantly during the first month. For the first two days, proportions are near the one- to twenty-year average of 60.59%. Values subsequently rise for about two weeks, abruptly decrease around the twentieth day, and shortly return to standard level.

The myeloid-erythroid ratio is low approximately two days but reaches 11.18 by the end of the second week. During the year, the value slowly descends to 2.95.

Individual erythroid and mycloid types, especially large subgroups such as polychromatic normoblasts and metamyelocytes, follow the trends of their own series.

Lymphocytes are few for a week and increase to 40% around the twentieth day. Throughout the first and second year the level declines, corresponding to the decrease in peripheral blood, to a stable average of 16.03% at the age of 2 or 3 years.

The relatively scanty numbers of basophilic granulocytes, plasma cells, megakaryocytes, monocytes, and histiocytes do not alter with age.

#### Penicillin Aerosol Inhalation for Babies

P. KARLBERG, M.D., G. STERNER, M.D., AND G. WALLMARK, M.D.<sup>‡</sup>

FOR penicillin inhalation by infants and small children, a small plexiglass dome, edged with plastic tissue, is very convenient.

Treatment is especially useful for acute respiratory infection of patients too young for a mask and for chronic illness requiring prolonged medication, such as bronchitis or nephritis.

P. Karlberg, M.D., G. Sterner, M.D., and G. Wallmark, M.D., treated 30 children aged one month to two and a half years at the Norrtull Hospital, Stockholm. A ten-minute inhalation provided the same blood level as an equal amount injected intramuscularly.

The dome, which covers the baby's head, is supplied in two sizes, holding either 6.5 or 10.5 liters. The soft plastic tissue is draped around the child's neck and upper chest, and aerosol is blown in from the back rather than over the face. Even struggling patients soon grow calm.


Crystalline sodium penicillin G in 2 cc. of physiologic saline solution is vaporized by Barach's spraying device driven by oxygen from a cylinder. Children up to 20 kg. in weight are given 50,000 units at a time and heavier patients as much as 100,000 units. From three to six inhalations are administered daily.

In severe, disease, aerosol is supplemented with intramuscular or oral penicillin or sulfonamides.

Aerosol-penicillin therapy for infants and children. Ann. paediat. 175:263-273, 1950.

# When You Consider

the High Cost of Babies .



That's why so many doctors everywhere are recommending Pet Evaporated Milk for infant feeding! Pet Milk does assure babies of optimal nutrition... does give parents the benefits of maximal economy!

You know that easy-to-digest Pet Milk is uniformly rich in all the food values the best milk can be depended upon to supply! You know that it is surely safe... because Pet Milk is sterilized in its sealed container!

Yet Pet Milk, the original evaporated milk, costs less than any other form of whole milk—far less than special infant feeding preparations!

So recommend Pet Milk for the babies in your care! It's nutritious! It's safe! And it keeps the cost down, too!

Favored Form of Milk for Infant Formula



PET MILK COMPANY, 1484-B Arcade Bldg., St. Louis 1, Mo.

§ EXTRARENAL AZOTEMIA and postrenal or obstructive uropathy can usually be differentiated from primary renal azotemia by the disproportionately low blood creatinine, compared with the elevated nonprotein nitrogen range. Although creatinine may be relatively high in all three conditions, no consistent relationship with nonprotein nitrogen can be traced. The two values were studied in 44 cases of urinary disease by Peter Gaberman, M.D., and associates at Cook County and Mount Sinai hospitals, Chicago.

#### Management of Postoperative Urinary Retention

VERNON S. DICK, M.D.\*

INABILITY to urinate after operation may be caused by acute bladder decompensation from rapid fluid intake, general sensory depression, prostatic or urethral obstruction, or combinations of these and related factors.

The postoperative bladder should not be allowed to become overdistended. If other means fail, catheterization should ordinarily be performed before distention has exceeded 450 cc.; a palpable bladder usually indicates 350 cc.


Factors influencing urinary retention include the position of the patient; the amount, type, and rate of injection of intravenous and other fluids during and immediately after surgery; fluid loss; and the degree of narcosis.

Bladder sensation is usually diminished in the still partially anesthetized or narcotized patient, and excessive urine may collect before pain is felt. Also, if the operative site is so located that pain is produced by effort to contract the voluntary abdominal and perineal muscles, vesical emptying is inhibited.

As most adults have difficulty voiding when lying in bed, Vernon S. Dick, M.D., of the Lahey Clinic, Boston, suggests that patients who are to remain supine after major operations should practice urinating in the recumbent position for a few days before surgery.

The frequency of catheterization must be related to intake and to loss of fluid by sweating, respiration, bowel discharges, and other means of elimination. For patients who tolerate intermittent catheterization poorly or when circumstances indicate considerable delay before resumption of micturition, an indwelling catheter is advisable. A urinary antiseptic should be given to patients being catheterized intermittently or with inlying catheters.

# Management of postoperative urinary retention. Lahey Clin. Bull. 7:57-62, 1950.



Unit Packaging Headquarters presents The Moisture Proof Package of the Month

You cannot take for granted the condition of an antibiatic - the odds are too great, the autome loa grave. Sharpe and Dahme recognized their responsibility and asked [4, for the right package.

As you know, the potency of antibiotics thisappears in the presence of maisture. The Unit Package is not just a convenient carrier for labell or triurate but is intal learned in the effectiveness of the drug. The package must be absolutely maisture proof. Samples of every botch of Pencillin when packaged, are tested by the Government before being released for sale because the Perfection of the Package can spell life or death for the patient.

This I.L. Unit is made of a parous proof metalized combination of materials. To assure absolute moisture proof protection, the righter is sealed in a circular packet for only in a found center section dans the material be uniformly drawn about a round tablet and the flange sealed without a winkle.

Since inspection means Perfection a production parkage is submerged in water under vocuum presurfe every ten minutes to these the moisture protection of this reperb 11. Package.

Success or failure of treatment may after be dependent on the parkaging of the product. Your confidence in and preference for drugs which are 11. Unit Paskaged will be amply rewarded.



The Finest Packaging In The World to Live to

IVERS-LEE COMPANY

### Medical Forum

Discussion of articles published in Modern Medicine is always welcome. Address all communications to The Editors of Modern Medicine, 84 South 10th St., Minneapolis 3, Minn.

#### Surgery of Acute Cholecystitis

to the editors: There is no question, as Dr. Frank Glenn says, that cholecystectomy is the treatment of choice in acute cholecystitis if the patient has been properly prepared and the operation is not contraindicated by pathology in other systems.

The difficulties of cholecystectomy in acute cholecystitis are encountered technically when adequate surgical help and facilities for postoperative care are not available. For this reason, cholecystostomy should never be discarded as a method of therapy in these cases, particularly when surgery is undertaken in rural communities where there is not the help or facilities to care for these patients. The repair of damaged common ducts is difficult in the most experienced hands; in the hands of less experienced operators, the results are not good and may cause death.

The following is an excerpt from the conclusions of an article (S. Clin. North America 25:285:300, 1945) concerning a group of 332 cholecystitis patients whom I studied intensively:

 Acute cholecystitis is a progressive disease in which the complications of gangrene, perforation, and abscess may occur in the presence of subsiding clinical symptoms, signs, and laboratory

\*Modern Medicine, Nov. 1, 1950, p. 89.



data. These complications increase the mortality and morbidity of the disease.

● In this series, the mortality and morbidity in acute cholecystitis were less when cholecystectomy was performed before the complications occurred.

 Cholecystectomy in the presence of the three main complications requires careful analysis of its indications in the individual case.

 Cholecystectomy for acute cholecystitis is safer when the gallbladder is removed from the fundus down.

 Cholecystostomy is not the operation of choice in acute cholecystitis unless local or systemic disease contraindicates removal of the viscus.

◆ The best results from cholecystectomy for acute cholecystitis depend upon the active cooperation of the medical men who first see the case.

Individual pathologic, local, and systemic response to gallbladder infection



# Veratrite®

In Mild and Moderate Hypertension, a marked sense of well-being is provided by the routine administration of Tabules Veratrite. Pronounced relief of headache, dizziness, fatigue and nervous irritability is accompanied with a reduction in blood pressure. Veratrite lessens peripheral resistance without compromise of circulation.

**Supplied:** Bottles of 100, 500, 1000.

SAMPLES AND LITERATURE ON REQUEST.

IRWIN, NEISLER & COMPANY



DECATUR, ILLINOIS

in middle-aged to elderly persons does not have such uniform pattern that a simple surgical therapeutic rule can be applied to the age group. Treatment must be individually rationalized.

BEVERLY CHEW SMITH, M.D.

New York City

► TO THE EDITORS: The removal of gallstones as treatment for acute cholecystitis is now an established fact.

Although surgical mortality is low for elective operations on the biliary tract, it becomes considerably higher when complications such as acute cholecystitis or common duct stone occur.

Acute gallbladder disease has been repeatedly shown to be due to an occlusion of the cystic duct by a stone followed by the typical clinical and pathologic effects produced by subsequent chemical changes. Infection may be present but generally plays a secondary role to the occluded duct.

Perforation of the gallbladder with peritonitis is a serious complication of acute cholecystitis and may result from a gangrenous area in the fundus or through a Rokitansky-Aschoff sinus. This complication often occurs with minimum signs and symptoms. It cannot be overemphasized that the clinical and laboratory findings in acute cholecystitis do not always indicate the severity and progression of the underlying pathologic process.

The so-called "silent gallstone" is a potential source of cystic duct occlusion and should be removed. Procrastination may end in a fatality because the patient has not been impressed with the importance of silent stones. Frequently, such stones produce severe reflex cardiac symptoms resulting in years of unnecessary treatment and restrictions for alleged heart disease which could easily be eradicated by elective operation.

More thought should be given to the preventive aspects of acute cholecystitis by advising surgery immediately after gallbladder disease is discovered, regardless of the patient's age.

The risk of surgery as a preventive measure in the early and mild forms of the disease is far less than in the later complications of acute cholecystitis, perforation with peritonitis, and common duct stone.

In a population of increasing longevity this principle is assuming greater importance since it is in the elderly that the incidence of gallstones reaches its peak. Every physician should review all aspects of his management of gallbladder disease and weigh the risks of surgery in the early stages as a preventive measure against a probable higher mortality rate in later years.

H. T. GROSS, M.D.

Appleton, Wis.

TO THE EDITORS: Whether an acute gallbladder should be removed or even drained is still a debatable question. Surgical judgment is difficult to evaluate, because each case presents individual problems.

As a general rule, subject to exceptions, I would say that acute obstruction of the cystic duct with associated hydrops should have a chol-

# Subdue Gastrointestinal Revolts...with

# **EMETROL**

PHOSPHORATED CARBOHYDRATE SOLUTION

- · Physiologic-not pharmacologic-action
- Free of antihistaminics, barbiturates, narcotics, and stimulants
- · Nontoxic-no distressing side-effects
- · Works quickly
- · Very agreeable taste
- · Simple regimen

prevent
or
arrest
nausea
and
vomiting
of
gastrointestinal
origin
quickly..
safely..
simply..

Clinical findings<sup>1</sup> in a series of 243 cases have demonstrated the value of EMETROL\* in preventing or arresting nausea and vomiting of nonorganic origin. A few doses usually suffice to overcome epidemic vomiting ("intestinal flu") in children or grownups; regurgitation in infants; and motion sickness. Many physicians are also employing EMETROL to check the nausea and vomiting of early pregnancy, as well as the nausea accompanying the administration of certain antibiotics.

SUPPLIED: Bottles of 3 fl.oz. and 1 pt., at all prescription pharmacies.

Bradley, J. E.: To be published.
 Trademark of Kinney & Company.

For detailed information on action, desage, and administration, send for complete literature.



#### KINNEY & COMPANY

Prescription Products

COLUMBUS

#### Salvage Possibilities in Threatened Abortion\*

TO THE EDITORS: What is the preferred treatment of threatened abortion? This question was recently raised in the Medical Forum of Modern Medicine, using the article, "Salvage Possibilities in Threatened Abortion," by Drs. Emmett D. Colvin, Rudolph A. Bartholomew, William H. Grimes, and John S. Fish as basis for the discussion. As these authors point out, very few have approached the subject from the clinical angle and the salvage possibilities, although Stander did so in 1942 in a study in which I collaborated (Am. J. Obst. & Gynec. 44:531, 1942).

Dr. Colvin et al. present a statistical study of 1,570 private cases which is postpartum and postabortal in its scope and, therefore, has the advantage of hindsight. Patients with very early pregnancies were observed at home, while those more advanced were hospitalized.

At present, we hospitalize virtually all these patients, using bed rest, sedation, plasma vitamin C determinations, cultures, and a thorough investigation to determine various factors, defects, and deficiencies as possible causes of abortion. At times, thyroid extract, ascorbic acid orally and intravenously, and vitamin K are administered as indicated. Coitus is interdicted for the remainder of the pregnancy. The use of mineral oil is also forbidden.

If the pregnancy continues, well and good. Otherwise, much clinical, laboratory, and pathologic data have

\*Modern Medicine, Oct. 15, p. 91.

been obtained during the hospital stay which will be of considerable value in the prognosis and successful management of a subsequent pregnancy.

Dr. Colvin and associates' results can be divided into three major groups: 1,098 going to term, 440 having abortions, and 32 having premature labor. These patients had received prenatal care and routine multivitamins, the exact nature and dosage of which were not revealed. In recent years, vitamin C has been added to these preparations in amounts ranging from 20 to 180 mg. per capsule. The vitamin C may have influenced their good results, since the impressive percentage of fullterm infants of 69.9% is somewhat better than the 60.5% reported by Stander without specific therapy. The authors do not provide control data to show that multivitamins did or did not influence the results.

It is difficult to reconcile the authors' statement that the prophylactic use of various hormones and vitamins in threatened, spontaneous, and habitual abortion "is not justified" with their actual use of "the usual multivitamins routinely administered during antepartum care." The fact that 69.9% continued to term is not restated in the summary and conclusions. Instead, they state, "The general assertion and acceptance of the prophylactic value of hormone and vitamin therapy in threatened or habitual abortion are apparently unwarranted."

There were 440 cases, or 28%, ending in abortion. It was suggested that vitamin and hormone therapy would have been given to these

no Querhose Muissan

with full-footed

# ACE Elastic Hosiery

here's why ...

Differing from the usual elastic stocking,

ACE® Elastic Hosiery is knit with a full
nylon toe... is form-fitted... is as
sheer in appearance as service-weight
nylons... and is offered in an
inconspicuous neutral shade of beige tha
blends with the smartest apparel.

Elimination of overhose does away with the unattractive bulkiness, uncomfortable weight, and unsightly wrinkles which have made patients rebel against supportive stockings.

WOMEN WILL WEAR

WITHOUT EMBARRASSMENT

AND IN COMFORT!

#### suspension support

The full foot in ACE Elastic Hosiery gives positive terminal anchorage immediately behind the toes. When the hose is drawn on the leg under tension and fastened with garters, it provides a radically new type, sheath-like supension support to venous structures of the leg.

#### fitted by "Calcufit"

ACE Elastic Hosiery is quickly and easily fitted with exclusive B-D "Calcufit" Charts supplied free to physicians.

Fashioned by B·D
akers of ACE elastic bandages
Chers of Mos classic manages

B-O PRODUCTS

Made for the Profession

since 1897

Picase sen		O COMPANY,		
Dr.	LASTIC MOSII			
Street			100	

while the abortion was in progress and, therefore, would be of little value. The proponents of such a therapy do not disagree with this point of view, since it is generally agreed that it is preferable to begin prophylactic therapy before the patient is pregnant, especially with a history of previous threatened, spontaneous, or habitual abortion. These patients need special detection and correction of as many of the contributory factors, defects, and deficiencies as possible, in order to be managed individually in subsequent pregnancies. Statistical summaries of end results do not help in the immediate diagnostic and therapeutic management of an actual patient with threatened abortion. New therapeutic agents must be constantly employed and investigated in an attempt to lower the large percentage of abortions.

Of the 440 abortions, 318 or 72% were classified as blighted ova, one of which is illustrated in Figure 11 of their original article. This illustration also shows a large amount of decidual hemorrhage. In our experience, such bleeding, often shown to be old because of hemosiderin pigment in the decidual cells, is sometimes associated with blighted ova. Such environmental factors were suspected by Mall and Meyer as the cause of some of the abnormal ova in their series, but they lacked the data. Both placenta praevia and abruptio placenta have decidual bleeding and these lesions had a combined incidence of 16% in their group of 60 cases with fetal abnormality.

It would have been of interest to

know the incidence of placenta previa and premature separation of the placenta in the entire series of 1.570 cases. Stander found an incidence of placenta previa at term of 2.9% in his study of 489 patients with threatened abortion. In our preliminary study of 500 cases of spontaneous abortion (Texas State 1. Med. 46:780, 1950), decidual hemorrhage had an incidence of 45%, and placenta previa of 4%. Prophylactic therapy with fresh orange juice and vitamin C supplements may be of value in controlling decidual hemorrhage. Such a nutritional regimen of adequacy. including the correction of all abnormal factors, defects, and deficiencies, has yielded 80% full-term pregnancies in a number of habitual abortion patients (Am. 1. Obst. & Gynec. 57:878, 1949).

Dr. Colvin et al. raise the question, "Is it not objectionable to administer any treatment which will dull the irritability of the uterus, and therefore permit prolonged retention of a product which has dangerous potentialities [i. e., malignant mole]?" It seems logical to cope with this fear by performing more operative completions of the abortion to provide the pathologist with additional tissue for study. Their incidence of curettage was only 3.4%.

In the discussion of this paper, Dr. Seeley advocated more curettements. Greenhill advocates frequent curettage for incomplete abortion. Stander found that 67% of 1,862 cases required operative completion.

The chief indication for intervention is profuse continued bleeding. Septic cases should not be invaded



"a unanimous decision"

# Eskadiamer tastes good!

Children—and adults—actually like to take Eskadiamer because it is so unusually pleasant tasting . . . and because it is not thick and cloving but light and easy to swallow.

Eskadiamer contains equal parts of sulfadiazine and sulfamerazine—the two safest sulfonamides in general use.

Since Eskadiamer is an aqueous suspension of microcrystalline sulfadiazine and microcrystalline sulfamerazine, desired serum levels are attained much more rapidly than with sulfonamide tablets of equivalent dosage.

Smith, Kline & French Laboratories, Philadelphia

# Eskadiamer the delicious fluid preparation

of sulfamerazine and sulfadiazine

'Eskadiamer' T.M. Reg. U.S. Pat. Off.

Each 5 cc. (one teaspoonful) of Eskadiamer contains 0.25 Gm. (3.86 gr.) microcrystalline sulfamerazine and 0.25 Gm. (3.86 gr.) microcrystalline sulfadiazine—the dosage equivalent of the standard 0.5 Gm. (7.7 gr.) sulfonamide tablet.

unless hemorrhage forces the issue. In these, a preliminary course of penicillin and sulfadiazine for several days may render the completion less hazardous. Intravenous pitocin drip also has been employed successfully in a few cases with the curettage. It greatly reduces blood loss during the procedure.

CARL T. JAVERT, M.D.

New York City

#### Fractures of the Hand\*

TO THE EDITORS: Dr. E. James Morrissey in his paper on fractures of the hand has dealt with the importance of these fractures and the frequency with which they are brushed aside as minor injuries. Hand fractures are tremendously important.

Very few fractures require pulp traction. It is very important, in splinting, to place the hand in physiologic rest, and this is thoroughly

dealt with in the paper.

With chip fractures of the distal tuft of the distal phalanx, it has been my experience that very few patients have any disability. In the baseball finger, another method of fixation is to flex the finger into the base of the hand and to hyperextend the distal joint by a thick pad of gauze under the end of the finger, keeping it strapped in this position for six weeks.

For fractures of the metacarpals, it is my belief that fixation by Kirschner wire is the method of choice and produces the best results at the earliest time. This applies to fractures of the neck or shaft and is not necessary for most fractures of the base. \*MODERN MEDICINE, May 15, 1950, p. 66.

The principle is that the fracture is reduced by traction and countertraction.

If the second metacarpal is broken, the wire is passed laterally through the head of the second into the third and is left protruding about 1/2 in. through the skin. While traction is maintained, a Colles type of plaster is applied around a cork over the end of the wire. By this method there are two fixed pointsthe head of the third metacarpal, and the cork in plaster-and the fracture cannot change position.

Fractures of the third are dealt with in the same way. Fractures of the fourth and fifth are wired from the ulnar side of the hand in a similar fashion. This permits full finger movement within a few days

and a very useful hand.

In fracture dislocation of the base of the first metacarpal, Bennett's, I feel that the most satisfactory method of fixation is by a Kirschner wire. The dislocation is reduced by traction on the thumb and index finger in the position of grasping, so that index and thumb could touch. When reduction is perfect, a Kirschner wire is passed through the distal part of the first metacarpal and into the shaft of the second metacarpal, and a cork is placed over the projecting 1/2 in. of wire. A plaster cast is applied to include the cork, up to, but not including the distal joint of the thumb.

The advantage of this method is that fixation is adequate and the patient can grasp objects between thumb and index finger.

H. R. C. NORMAN, M.D.

#### THE NAME TO DEPEND ON-

# DEVILBISS

A comprehensive development and research program assures you of the most efficient in nebulizers and atomizers



The continuing DeVilbiss research program, now more comprehensive than ever before, is keeping pace with the latest findings in the field of aerosol therapy.

DeVilbiss' close cooperation, particularly with the pharmaceutical houses and research foundations, has resulted in the exactly *right* piece of equipment being

always available for new techniques. There is an atomizer available for any new combination of pH, viscosity, or surface tension likely to be encountered.

DeVilbiss deserves your confidence. Over the years, no other manufacturer of similar equipment has worked so closely with research development to the end that the physician would have exactly the right type of equipment for the job at hand.

De Vilbiss...

ATOMIZERS NEBULIZERS VAPORIZERS

## Diagnostix

Here are diagnostic challenges presented as they confront the consultant from the first clue to the pathologic report. Diagnosis from the Clue requires unusual acumen and luck; from Part II, perspicacity; from Part III, discernment.

#### Case MM-185

#### THE CLUE

like to discuss with you is a 20-yearold girl. She has bilateral sciatica which becomes worse with exertion and is aggravated by coughing and sneezing. She has had dull pain in the left sacroiliac and



left gluteal region every night for the past month and gets relief by rubbing with liniments and from heat and aspirin. The condition is of two years' duration and, although she is incapacitated now, she is better than a year ago.

VISITING M.D. What's the history?

ATTENDING M.D. Two years ago she began to have catches in her back so that she could not straighten up after leaning forward and lifting anything. These occurred inabout termittently for months, at which time the leg muscles began to ache. A murmur was heard over the heart, and some twitching choreiform movements were observed. This illness was diagnosed as rheumatic fever by the attending physician and she was hospitalized for three months. During this time the left sciatic pain began and became quite severe. When permitted to get up she walked with a limp because of the pain. She had lost an inch and a half in height and the doctor noted definite kyphoscoliosis.

#### PART II

VISITING M.D: Was there any effect from strain at that time?

aftending M.D. Coughing and sneezing aggravated the pain in the left sacroiliac region and sciatic radiation. The discomfort from straining at stool was sufficiently severe so that on occasion she resorted to enemas or laxatives. There was some tenderness over the third and fourth lumbar vertebrae, on the left more than on



# HEXACHLOROPHENE SOAP

As an Adjunct In Treating Pyogenic Infections

As a Skin Antiseptic in Your Office and Home

## **GAMOPHEN**

(HEXACHLOROPHENE)

Surgical Soap

NOTHER ETHICON QUALITY PRODUCT

ETHICON SUTURE LABORATORIES, INC.

NEW BRUNSWICK, NEW JERSEY

Gamophen Hexachlorophene Surgical Soap has been accepted by the Council on Pharmacy and Chemistry of the American Medical Association. It is a highly purified detergent soap containing hexachlorophene, an efficient antiseptic that adds to the cleansing action of the soap a bactericidal and cumulative bacteriostatic effect, even in high dilution.

Because of its prolonged antibacterial suppressive action, it has been found not only to be an excellent agent for preparing the hands of the surgeon in the operating room, but also a valuable adjunct in the treatment of certain skin conditions involving pusforming infections.

#### FREE-FULL-SIZE BAR FOR TRIAL

(May be clipped and passed to Penny Post Card)
ETHICON, New Brunswick, N. J. DEPT. MM-251
Please send Gamophen Soap and Literature.

Dr.\_\_\_

Street

 the right. Five months ago the sciatic radiation appeared on the right. About four weeks ago a dull constant pain kept her awake two nights.

visiting M.D.: Was she unable to sleep because of pain, or did the pain waken her out of a sound sleep?

ATTENDING M.D. She was wakened by sudden pain and could not sleep for the following forty-eight hours.

VISITING M.B: Were there any important clues in the story you obtained from her?

ATTENDING M.D: No. She had no illness or sciatica before the present episode. Two years ago a classmate who sat across the aisle from the patient died of tuberculosis. She was coughing and spreading the tubercle bacilli and 2 other classmates died of the disease. visiting M.D: What do the chest and spine roentgenograms show, and what is the tuberculin test?

ATTENDING M.D: The chest film shows kyphoscoliosis and some fuzziness of the outlines of the left third and fourth lumbar foramina, which seem a little wide. There may be erosion of these vertebrae posteriorly. The tuberculin skin test was negative.

VISITING M.D: How did the radiologist interpret this?

ATTENDING M.D: He called the film negative, saying the widening of the foramina was inconclusive.

#### PART III

visiting M.D.: We are concerned here with a young person with kyphoscoliosis which she did not have before the recent illness. The situ-

(Continued on page 154)



"Remind me, sergeant, to write an article blasting socialized medicine, will you?"

#### Doctor . . .

Here are two great Spot Tests that simplify Urinalysis.

### GALATEST

The simplest, fastest urine sugar test known.

### **ACETONE TEST**

(DENCO)

For the rapid detection of Acetone in urine or in blood plasma.

A LITTLE POWDER
A LITTLE URINE



COLOR REACTION IMMEDIATELY



Combination Kit: Contains both tests, a dropper and color chart. Available at all drugstores and surgical supply houses.

#### BIBLIOGRAPHY

Joslin, E. P., et al: Treatment of Diabetes Mellitus—8 Ed., Phila., Lea & Febiger, 1946—P. 241, 247.

Duncan, G. G., Carey, L. S. & Hudson, M. T.: Medical Clinics of North America—Phila., No., W. B. Saunders, Co. Phila., 1949—P. 1538.

Lowsley, O. S. & Kirwin, T. J.: Clinical Urology—Vol. 1, 2 Ed., Bult., Williams & Wilkins, 1944—P. 31.

Duncan, G. G.: Diseases of Metabolism— 2 Ed., Phila., W. B. Saunders Co., 1947—P. 735, 736, 737.

Guidotti, F. P. & Winer, J. H.: The Military Surgeon-Vol. 94, Feb., 1944, No. 2.

Bacon, Melvin: The Journal of The Maine Medical Assn.—Vol. 38, July, 1947, No. 7—P. 167. Stanley, Phyllis: The American Journal

Stanley, Phyllis: The American Journal of Medical Technology-Vol. 6, No. 6, Nov., 1940 and Vol. 9, No. 1, Jan., 1943.

#### GALATEST and ACETONE TEST (Denco)

... Spot Tests that require no special laboratory equipment, liquid reagents, or external sources of heat. One or two drops of the specimen to be tested are dropped upon a little of the powder and a color reaction occurs immediately if acetone or reducing sugar is present. False positive reactions do not occur. Because of the simple technique required, error resulting from faulty procedure is eliminated. Both tests are ideally suited for office use, laboratory, bedside, and "mass-testing." Millions of individual tests for urine sugar were carried out in Armed Forces induction and separation centers, and in Diabetes Detection Drives.

The speed, accuracy and economy of Galatest and Acetone Test (Denco) have been well established. Diabetics are easily taught the simple technique. Acetone Test (Denco) may also be used for the detection of blood plasma acetone.

Write for descriptive literature

THE DENVER CHEMICAL MFG. CO., Inc. Dept. 2, 163 Varick Street, New York 13, N. Y.

# without distraction

# · Cardiac Decompensation


# in Coronary Disease

# in Hypertension

The superior features of theobromine therapy are now available without the side-effects which drive doctors to distraction. Maltbie research has produced an unique double salt of calcium theobromine and calcium gluconate-











#### Seeing is believing

Calpurate is really different from mere mechanical mixtures of theobromine and calcium gluconate. Convincing evidence may be seen at a glance in the accompanying polarized photomicrographs. Compare the rectangular crystals of theobromine (top), and the needle-like crystals of calcium gluconate (middle) as found in mixtures with the double-salt crystals from a solution of Calpurate (bottom).

#### In clinical practice, too

Seeing is believing...

Calpurate affords sustained myocardial stimulation, vasodilation and diuresis without gastric disturbances.
Calpurate with Phenobarbital relieves stress and improves circulatory efficiency, exerts a desired calming effect.

Administration and Dosage: 1 or 2 tablets t.i.d.

Maltbie Laboratories, Inc., Newark 1, New Jersey



# Calpurate 'Tablets and Powder

Theobromine Calcium Gluconate Maltbie

# **Calpurate**

with Phenobarbital . Tablets

for trouble-free, prolonged cardiac therapy

ation is confused by two red herrings: the diagnosis of chorea and rheumatic fever, and the tuberculous roommate. Of course. one thinks at once of a ruptured disk, but there are several inconsistencies. Neurologic examination is completely negative, there is curvature of the spine, atrophy of the left leg, some loss of weight and height, and percussion tenderness over lumbar 3, 4, and 5. I find Lasègue's sign negative and ankle jerks and sensation normal. The patient has an intraspinal lesion and I think we should immediately have a myelogram. I believe this is probably a tumor, not a disk. There is no evidence of trauma, and the condition is progressive despite the fact that she has less pain than a year ago.

#### PART IV

surgeon: (The myelogram revealed a suggestive lesion at lumbar 2, and surgical exploration is being done.) We're doing a bilateral lumbar laminectomy. An enormous tumor extends from the conus to the fourth lumbar vertebra. The growth is not well incapsulated, and many of the nerve roots are adherent to the capsule. strange that with this much infiltration in the nerve roots the patient did not have more nerve root symptoms. The tumor seems to arise from a filum terminale. The lower end of the tumor is about lumbar 4, which is where the sciatic nerve begins.

PATHOLOGIST: (After examining the tumor) This tumor is an ependymoma. The prognosis is fair.



"I'm the one who's supposed to say 'ah'!"

#### GRATIFYING IMPROVEMENT IN ACNE, KELOIDS, WITH NEWLY ISOLATED

ISOLATED LIVER FRACTION



IN ACNE

reduces papules, pustules, comedones<sup>1,2,3,4</sup> • modifies unsightly scarred areas<sup>1</sup>

IN KELOIDS

reduces size of keloids<sup>4</sup>
• inhibits regrowth after surgical removal<sup>5</sup>

KUTAPRESSIN\* restores normal skin tone in acne—by constricting capillaries, decreasing capillary volume, and increasing the rate of blood flow through affected tissues.<sup>1,4</sup> This eliminates the passive hyperemia arising from local stasis of blood and tissue fluids, and restores the skin's resistance to secondary infection.

In keloids, constriction of capillaries and reduction of their permeability prevents flow of blood serum into subcutaneous tissues, thus decreasing the distention of the skin and the accompanying inflammation. 4.6 Administration of **KUTAPRESSIN** before, during, and after surgical removal of keloids decreases loss of blood serum into the site of scar formation and inhibits regrowth. 5

REFERENCES: 1. Marshall, W.; J. M. A. Alabama 13: 255 (1941). 2. Lichtenstein, M. R., and Stillions, A. W.; Arch. Dermat. & Syph. 45: 959 (1942). 3. Stillions, A. W.; Missistippi Valley M. J. 64: 135 (1942). 4. Marshall, W., and Schadeberg, W.; Wisconsin M. J. 49; 369 (1950). 5. Marshall, W.; Paper read before Midwestern Section of the American Federation for Clinical Research, November 2, 1950.



Available in 10-cc. multiple-dose vials through your usual source of supply.

\*\*Supply: \*\*Supply

<sup>\*</sup>Trademork of Kremers-Urban Co.

#### Basic Science Briefs

Obstetrics

#### **Blood Volume in Pregnancy**

During gestation, the volume of the mother's heart increases about 170 cc., and blood volume about 1,500 cc. Dr. S. R. Kjellberg and associates of Karolinska Sjukhuset, Stockholm, conclude that the heart enlarges chiefly because of greater blood content but partly through myocardial growth. Variations in heart and blood volumes are not directly proportional. Heart contours of 23 primiparas and 23 multiparas aged 18 to 44 years were examined radiologically, and cardiac volume was calculated by a modification of the Larsson-Kjellberg technic. Blood voltime was estimated from relative and total hemoglobin, the latter determined by Sjöstrand's carbon monoxide method.

Acta med. Scandinav. 138:421-429, 1950.



"Is your name on the sick book? Stick out your tongue . . . Corporal give him an APC."

Metabolism

#### Pituitary Function

The decreased ability of hypophysectomized rats to synthesize glycogen from glucose is restored by ACTH. Greatly disturbed phosphorus levels also return to normal, report Drs. L. G. Abood and J. J. Kocsis of the University of Chicago and the Armour Research Laboratory, Chicago. Anaerobic glycolysis of the brain, after 20% reduction by pituitary resection, is likewise improved by the hormone.

Proc. Soc. Exper. Biol. & Med. 75:55-58, 1950.

Cardiology

#### Saliva Electrolytes

Congestive heart failure is associated with unusually low concentrations of sodium and chloride in the saliva and with high potassium levels. No difference is noted whether the diet is high or low in sodium, with salt intake ranging from 1 to 10 gm. or more daily. In contrast, noncardiac subjects on a low-salt regimen have less salivary sodium and chloride than a group receiving a regular diet, and electrolytes are in the range for cardiac patients. Dr. Abraham G. White and associates of the Montefiore Hospital, New York City, found no correlation between electrolyte concentrations of saliva and blood. Samples were collected in the morning before the patients had

J. Clin. Investigation 29:1445-1447, 1950.

# Today's trend is to liquid oral penicillin

"...it has been demonstrated repeatedly that the oral route is as effective as the parenteral route when adequate doses of penicillin are used."

Keefer, Chester S.: Am. J. Med. 7:216

# Eskacillin Eskacillin

The unusually palatable liquid penicillins for oral use

In keeping with today's trend to oral penicillin, S.K.F. now offers, for your convenience, Eskacillin in 2 strengths: 'Eskacillin 100', containing 100,000 units of penicillin per 5 cc. (one teaspoonful). 'Eskacillin 50', containing 50,000 units of penicillin per 5 cc. (one teaspoonful).

#### Among the many indications are:

Acute sinusitis Pneumonia
Bronchitis Cellulitis
Tonsillitis Gonorrhea

Otitis media Certain skin infections

Smith, Kline & French Laboratories, Philadelphia

'Eskacillin' T.M. Reg. U.S. Pat. Off.

#### Nutrition

#### Sodium Deficit with Tuberculosis

Far advanced tuberculosis may be associated with an unusual type of sodium depletion causing no symptoms. Hyponatremia occurs without dehydration, peripheral circulatory failure, or hyperalkalemia, and adrenal and kidney function are seemingly unimpaired. Renal clearance of creatinine, mannitol, and urea is normal. Dr. E. A. H. Sims and collaborators of Yale University, New Haven. Conn., believe that the primary factor is reduction of cellular osmotic pressure owing to severe debility and malnutrition. blood sodium was observed in q patients with pulmonary tuberculosis and 1 with miliary infection. Values were normal, however, for 14 others with moderate or severe tuberculosis and in 5 instances of malnutrition associated with other disease.

I. Clin. Investigation 29:1545-1557, 1950.

#### Hypertension

#### Vasopressor in Blood

Pherentasin, a powerful pressor substance rarely found in healthy subjects, has been obtained in fairly pure form from arterial blood of hypertensive patients. The usual concentration is 10 to 20 y per liter. report Drs. Henry A. Schroeder and Norman S. Olsen of Washington University, St. Louis. When the same level is produced in rats, diastolic pressure rises 20 to 40 mm. of mercury for thirty to sixty minutes. Pherentasin is consistently obtained from the blood of patients with renal hypertension and, in smaller

amounts, from those with neurogenic or endocrine forms, but almost none from malignant hypertensive individuals. The vasopressor is nonprotein and contains an amine and a carbonyl group.

J. Exper. Med. 92:545-559, 1950.

#### Oncology

#### Motile Cancer Cells

The metastatic capability of malignant tissue may be partly related to independent ameboid motion of single cells. Tissue tends to separate because of calcium deficiency, and detached cells may invade interstitial spaces of neighboring healthy areas. By means of cinephotomicrographs, Drs. H. T. Enterline and Dale Rex Coman of the University of Pennsylvania, Philadelphia, recorded activity of material from human breast cancer, renal cancer, mesenteric leiomyosarcoma, mouse fibrosarcoma, and other animal tumors. Average speed of human malignant cells was 0.3 to 1.4 µ per minute and the maximum rate o.q to 4.4 µ. Benign glandular epithelial cells from a cystic breast traveled about 0.3 µ per minute, with maximum rate of 1.2 microns.

Cancer 3:1033-1038, 1950.



# when eating for two"

... plenty of citrus fruits

Most obstetricians today insist that their mothers ingest plenty of vitamin C, particularly after the first trimester' (8 oz. citrus juice during pregnancy, 12 oz. while lactating)." When an adequate nutritional regimen (with particular reference to vitamin C) is followed throughout pregnancy, toxemia is reduced'-more babies are born normally and with a higher birth weight" -premature and still births are fewer -- and both maternal and infant health are improved postpartum.\* Most mothers enjoy the flavor of fresh Florida citrus fruits (so rich in vitamin C and containing other nutrients\*), as well as the energy pick-up provided by their easily assimilable fruit sugars.3

\*Citrus fruits - among the richest known sources of estamin C-also contain estamins A and B. readily assimilable natural fruit sugars, and other factors, such as iron, calcium, citrates and citric acid

FLORIDA CITRUS COMMISSION LAKELAND, FLORIDA

#### References:

Notronances:

1. Burke, B. S. and Stuart, H. C.: J.A.M.A., 137:119, 1948. 2. Burke, B. S. et al.: Am. J. Obet, 6 Crocc. 60:38, 1943. 3. Burke, B. S. et al.: J. Nutrition, 20:559, 1943. 4. Javert, C. T. and Finn, W. E. Texas State. J. Med., 40:755, 1940. 5. McLester, J. S. Nutrition and Diet in Health and Diesase, Saunders, Phila, 4th ed., 1944. 6. National Research Cauncil: "Recommended Food and Nutrition Board, Daily Allowances for Specific Nutrients," Wash. D. C., 1948. 7. People's League of Health: J. Lancet, 2:10, 1942.







#### Short Reports

Endocrinology

#### Nebulized Cortisone for Bacterial Pneumonia

Symptoms of pneumococcic pneumonia are greatly modified by inhalation of cortisone, although bacteria are not affected. At the Henry Ford Hospital, Detroit, an elderly man with type VII infection was relieved of pain within three hours after commencement of therapy, appetite returned in five, and high fever subsided in twelve. Drs. William H. Reeder and George S. Mackey administered an aerosol containing 5 to 8 mg. every thirty to sixty minutes for thirty-three hours. The feeling of well-being was not accompanied by much change in roentgen appearance of lesions. With reduction of dosage, the temperature rose, and when the drug was withheld. toxicity, cyanosis, dyspnea, and pain returned.

Dis. of Chest 18:528-534, 1950.

Public Health

#### U.S. Cancer Facilities

The United States now contains 268 cancer detection centers, 165 cancer diagnostic clinics, 631 clinics for both diagnosis and treatment of malignancy, and 17 cancer hospitals. Services and facilities are listed for workers in the field in a source book recently published by the Federal Security Agency. The report was prepared by the National Cancer Institute of the Public Health Service.

Industrial Medicine

#### Women Absentees

In the industrial field, women leave their work 2.5 times as often as men for periods exceeding a week, reports Dr. W. M. Gafafer. Respective absences per 1,000 persons are 254.5 and 95.5, though men have a higher off-the-job accident rate, and major causes of ill health are about the same in the two sexes.

Antibiotics

#### Amebicidal Activity

For apparently the first time on record, an antibiotic has destroyed Endamoeba histolytica in vitro by direct attack rather than by inhibiting associated bacteria. Prodigiosin, a pigment extracted from surface cultures of Serratia marcescens, was employed in several dilutions against two strains by Drs. William Balamuth and Morgan M. Brent of Northwestern University, Evanston, Ill. At 1:2,000,000, individual protozoa became vacuolated, motionless, and pink and at higher concentrations were dissolved, with few persisting twelve hours. The NRS type of ameba grown with Aerobacter aerogenes was killed at 1:400,000, and even huge inocula of the more resistant UC strain associated with mixed bacteria were obliterated at 1:100,000. Related organisms, oxidation-reduction potentials, and pH were not adversely affected.

Proc. Soc. Exper. Biol. & Med. 75:874-878, 1950.



For maintaining the edema-free state, here—at last—is truly effective oral mercurial diuretic therapy. One or two Tablets MERCUHYDRIN® with Ascorbic Acid daily (more when indicated) generally controls cardiac edema with

greater convenience · greater economy · greater safety

# COMPERCUHYDRIN

After parenteral therapy, your patient has been brought to unfluctuating basic weight. Then systematic oral therapy employing Tablets MERCUHYDRIN (brand of meralluride) with Ascorbic Acid may eliminate the need for injections entirely in mild decompensation. In more advanced cases, you can greatly reduce the number of injections required to maintain your patients free of edema.

Prolongation of the interval between injections simplifies management. The diuretic response is good, the tablets are well tolerated, the method is convenient, and the economy considerable.

Packaging: Tablets MERCUHYDRIN with Ascorbic Acid, available in bottles of 100 tablets. Each tablet contains meralluride 60 mg. (equivalent to 19.5 mg. mercury) and ascorbic acid 100 mg.



#### Nutrition

#### Pancreatin for Avitaminosis

Vitamin A absorption is increased for patients with severe cystic fibrosis of the pancreas by administration of pancreatin. The glandular product employed by Dr. Gordon E. Gibbs of the University of Maryland, Baltimore, had lipolytic activity for 1 gm. equal to 40 cc. of duodenal aspirate, much higher than in material unsuccessfully used by others. In 32 cases, average plasma vitamin A mounted from 31 to 122 µg. three hours following doses of halibut liver oil. Without pancreatin, the rise was 34 to 37 µg. No consistent effect was produced for 10 patients with celiac disease, who had normal lipase before administration.

Pediatrics 6:593-600, 1950.



"But he's sick and I didn't want to call just any old doctor."

#### Nutrition

#### Alcoholism and Allergy

Chronic alcoholism is generally associated with sensitivity to native food ingredients of the favorite liquor, such as corn, wheat, grape, or apple. The pharmacologic response to repeated drinks is nonataxia, facial flush, and happy companionability, but the allergic response is ataxia, pallor, anorexia, nausea, vomiting, and morose, seclusive, and belligerent behavior. Dr. Theron G. Randolph of Chicago found masked allergy in 37 of 40 reformed alcoholics. Spirits induce and maintain chronic symptoms of sensitivity. Drinking the beverage immediately alleviates distress for the time being, and abstinence is followed by delayed increase in symptoms. Thus allergy may determine preference for specific liquors, compulsive drinking, and withdrawal effects. Proc. Central Soc. Clin. Research 23:84-86, 1950.

#### Dermatology

#### New Fungicide

An antibiotic principle of Streptomyces fradiae has in vitro activity against several aerobic actinomycetes and deep fungi. The term Neomycin C is suggested by Dr. Wilfred E. Wooldridge and Mary Hoffman of Washington University, St. Louis. Phialophora and several species of Nocardia were destroyed by the agent, although the dermatophytes and two yeast-like organisms were not affected. However, the fungicide may be toxic, since the lots of Neomycin containing the new factor are toxic to the kidneys.

1. Invest. Dermat. 15:351-353. 1950.

# Voctor... listen to this conversation about you!



PATIENT... There was so little discomfort in having that mole removed.

NURSE . . . The Doctor's new technic with the HYFRECATOR is wonderful!

PATIENT... You can tell the Doctor I'm going to have my sister come in. She worries about a mole on her abdomen.

NURSE ... It's a good idea to have him look at it to make sure ... and he'll know if it should be eradicated with the HYFRECATOR.

Over 70,000

#### **HYFRECATORS**

in daily use

to eradicate moles, warts, unwanted hair and other superficial growths. Many doctors use the HYFRECATOR...High Frequency Eradicator...for fulguration and bi-active coagulation as well as desiccation. They find that the HYFRECATOR'S double spark gap power, accurately controlled and smoothly graded current... exactly meets their individual office electrosurgical demands.

#### The BIRTCHER Corporation

Los Angeles 32, California

#### and Poctor ... Please Note

The increasingly widespread public knowledge of the danger of moles as fore-runners of skin cancer have made your patients more receptive to your suggestion that all suspect lesions be eradicated. New reprints on Precancerous Lesions are available. Write your name and address – mail for your copies.

# ERADICATE THAT SUSPECT PRECANCEROUS LESION

To: The BIRTCHER Corp., Dept. MM 2-51 5087 Huntington Dr., Los Angeles 32, Calif.

Please send me free reprints on Precancerous Lesions and Literature on the Hyfrecator.

Name.

Street.

Cirv

State

#### Medical Education

#### Poliomyelitis Fellowships

Candidates interested in viral, biochemical or biophysical, orthopedic, neurologic, epidemic, or other aspects of poliomyelitis may apply for pre- or postdoctoral fellowships extending one to three years, with the privilege of renewal. Stipends range from \$1,200 to \$1,800 a year plus tuition without doctorates, otherwise from \$3,600 to \$7,000. Applicants must be U.S. citizens in good health, with or working for an M.D., Ph.D., or equivalent degree. Marital and dependency status is considered in determining income. Complete information can be obtained from the Division of Professional Education. National Foundation for Infantile Paralysis, 120 Broadway, New York . N.Y.

#### Experimental Surgery

#### Pancreatic Duct Anastomosis

When the head of the pancreas is removed, the pancreatic duct should be sutured directly to an opening in the duodenal wall, with careful apposition and temporary use of a tube. Only a few more minutes are needed than for an implant or closure of the pancreas at the cut end. In determining the best procedure, Drs. Donald J. Ferguson and Owen H. Wangensteen of the University of Minnesota, Minneapolis, employed eight methods of pancreatic-intestinal juncture on 61 dogs: direct suture anastomosis with and without a tube, implantation without tube and with ordinary or very small tube, implantation by Varco technic, aseptic implantation

with necrotizing suture, and implantation of the sectioned end of pancreas. In most cases the duct was joined to the duodenum about 10 cm. below the original location.

Ann. Surg. 132:1066-1074, 1950.

#### Honors

#### Foreign Scientists Nominated

The New York Academy of Sciences nominated to foreign scientists as honorary life members at the 1950 annual meeting. The majority are in biologic fields, including Dr. William T. Astbury, English biophysicist; Dr. Charles H. Best, of the University of Toronto; Dr. Frank M. Burnet, Australian authority virus diseases; Dr. James B. Collip, medical dean at the University of Western Ontario; a Chilean biochemist, Dr. Eduardo Cruz-Coke; and Dr. Bernardo Houssay of Argentina. A research prize of \$200 was given to Dr. Paul B. Weisz, biologist at Brown University.

#### Medical Education

#### Clinical Chemistry Board

A national certifying board for those trained in clinical chemistry has been formed by representatives from three organizations: the American Chemical Society, the American Institute of Chemists, and the American Society of Biological Chemists. The new committee, analogous to the various medical specialty boards, is headed by Dr. Otto A. Bessey, University of Illinois, Chicago. Information may be obtained from the Secretary-Treasurer, Dr. J. W. E. Harrisson. 1921 Walnut St., Philadelphia.

#### FIBERGLAS\* REPORTS TO THE PROFESSIONS

# Fiberglas Cloth used as Backing for REESE DERMATAPE

(Reg. U. S. Pat. Off.,

(... A Skin Transfer Adhesive Tape)


A new technique for obtaining accurate skin grafts is made possible with the Reese Dermatape and the Reese Dermatomet. The technique enables any surgeon consistently and successfully to excise skin grafts from .008" to .034", to tailor the grafts accurately, and to transplant them without stretching or contracting, and usually without suturing.

An important feature of the Reese Dermatape is the coated backing cloth, woven of Fiberglas yarns.

This is important because:

- 1-Fiberglas cloth will not stretch.
- 2—Fiberglas cloth has enormous tensile strength, allowing it to be tightened on the Dermatome without danger of breaking.
- 3— Fiberglas cloth is impervious to aqueous and alcoholic solutions, permitting adequate sterilization by immersion.
- Fiberglas cloth securely anchors the rubber splint during excision of the skin graft.
- 5 Fiberglas cloth permits easy separation of the rubber splint and backing after excision of the graft, leaving the graft and splint intact for transplantation.

Inert, inorganic, nonallergenic, nonsensitizing and chemically stable Fiberglas fibers produce no harmful effect on human tissue . . . Owens-Corning Fiberglas Corporation, Dept. 29-B3, Todelo 1, Ohio.



†Developed by John D. Reese, M.D., Assistant Professor of Plastic Surgery, Jefferson Medical College, Philadelphia, Pa., in conjunction with Irvington Insulator & Varnish Co., Irvington, N. J., and with Lee Tire & Rubber Co., Conshohocken, Po. Dermatape is used with the Reese Dermatame, manufactured by Bard-Parker Co., Inc. (Agent), Danbury, Conn.



\*Fiberglas is the trade-mark (Reg. U. S. Pat. Off.) of Owens-Corning Fiberglas Corporation for a variety of products made of or with fibers of glass.

#### Endocrinology

#### Cortisone and Diabetes

The effect of adrenal hormones on rheumatoid arthritis is apparently not closely dependent upon metabolic influence. Patients with severe diabetes and active rheumatic disease were examined during various therapeutic regimens by Dr. E. M. Brown, Jr., and associates at the University of Pennsylvania, Philadelphia. To produce adrenal cortical response, the customary doses of insulin were reduced. Although urinary excretion of glucose, nitrogen, and ketones was increased, arthritis was not affected. When ACTH or cortisone was administered, the joint condition improved greatly, regardless of whether the manifestations of diabetes were exaggerated or controlled at the time.

J. Clin. Endocrinol. 10:1363-1374, 1950.

#### Pharmacology

#### Cocaine Toxicity

The amount of barbiturate usually given to prevent cocaine reactions is probably far too small. About half the anesthetic dose of pentobarbital, equal to 3 times the sedative dose, is required for protection of rabbits and dogs. In treatment of cocaine intoxication, only enough barbiturate is needed to stop convulsions. Larger amounts may cause respiratory failure, warn Drs. John E. Steinhaus and Arthur L. Tatum of the University of Wisconsin, Madison. Cardiovascular depression from therapeutic doses of cocaine may be counteracted by epinephrine. Contrary to report, the latter drug produced no ventricular fibrillation when given to 30 animals.

J. Pharmacol. & Exper. Therap. 100:351-361, 1950.



"It's most annoying to hear the interns refer to the medical technician as the urine Anna Liza."

#### Life's Weary Moments

Think of a gag that fits the illustration. For every issue a new gag is published and the author is sent \$5. The Feb. 15 winner is

A. H. Warner, M.D. Queens, N.Y. Mail your caption to The Cartoon Editor Caption Contest

No. 1 Modern Medicine 84 South 10th St. Minneapolis 3, Minn.

## Sturdy Soup For Babies – Heinz Strained Beef And Liver Soup

Infants Like The Flavor And Texture Of This Dish!



HEINZ makes a complete line of baby foods for your youngest patients! These quality products include —

Pre-Cooked Cereal
Pre-Cooked Oatmeal
Strained Baby Foods
Junior Baby Foods
— every one outstanding for
flavor, color and texture!


CHOICE beef, select livers, potatoes, carrots and other vegetables make Heinz Strained Beef and Liver Soup a wholesome, nourishing dish for the babies in your care! Expertly cooked and strained to a smooth consistency, this protein-rich soup is also a dependable source of riboflavin, iron and Vitamin A. And like all Heinz Baby Foods, Heinz Strained Beef and Liver Soup is made to a quality tradition mothers know and trust!

An 82-Year Reputation Backs the Complete Line of

## **Heinz Baby Foods**

CEREALS • MEATS • VEGETABLES
FRUITS • DESSERTS





#### Portrait of a former "coughing" patient

after his physician prescribed the highly palatable, non-narcotic Robitussin: distinguished by its intense and prolonged action in increasing respiratory tract fluid, and by its ability to improve mood.

(Glyceryl guaiacolate 100 mg., and desoxyephedrine hydrochloride 1 mg., in each 5 cc.)

### Robitussin<sup>®</sup>

is a product of A.H. ROBINS CO., INC. . RICHMOND 20, VA.

Ethical Pharmaceuticals of Merit since 1878



Appointments

#### New Director Is Named at National Institutes of Health

Dr. William H. Sebrell, Jr., succeeds Dr. Rolla E. Dyer as director of the National Institutes of Health under the Public Health Service. Formerly head of the Experimental Biology and Medicine Institute, the new director added greatly to knowledge of the vitamin B complex and was instrumental in drawing up the first international standards of nutrition for the League of Nations.

Public Health Rodent Control

Rats are outwitted by a type of poison, warfarin, that differs from all others in two respects: It kills only after repeated doses, and does not arouse enough suspicion for bait' refusal. Patent to the formula, 3-(alphaacetonylbenzyl)-4-hydroxycoumarin, is held by the Wisconsin Alumni Research Foundation. Rodents die of hemorrhagic shock caused by the inhibition of prothrombin formation and capillary injury. Drs. Wayland I. Hayes, Jr., and Thomas B. Gaines of the U.S. Public Health Service carried out successful field trials against wild Norway rats at Savannah, Ga., some in an old focus for typhus. From 85 to 100% of animals were killed at sites where ANTU was 50 to 80% effective and thallium sulfate 72 to 94%. Yellow corn meal containing 0.05 mg. per gram was an excellent bait. A total of 0.69 mg. per rat, or 2.87 per kilogram, was lethal, and infestation was generally controlled in two or three

weeks. Food was prepared with an electric mixer and dispensed in small glass bowls holding 300 cc. To protect other animals from eating the poisoned corn, bait may be placed in stations of wood and wire net. Presumably, dogs and cats eating the poisoned rats for several days may be killed.


Pub. Health Rep. 65:1537-1555, 1950.

Obstetrics
Antidiuretic Hormone

Blood from women in the last half of pregnancy has the ability to inactivate the antidiuretic action of the posterior pituitary hormone. The inhibiting factor, probably an enzyme, develops between the fourteenth and twenty-first weeks of gestation and gradually disappears after delivery. This antagonist of the antidiuretic hormone is employed by Dr. William I. Dieckmann and associates of the University of Chicago to evaluate the potency of posterior pituitary solution. Citrated blood of a pregnant woman and 0.04 unit of Pitressin are incubated for an hour at 100.4° F., and 10 cc. of plasma is withdrawn. Diuresis is established by 1,800 cc. of water swallowed in two and a half hours, and urine is obtained by catheter at thirty-minute intervals. After the first hour, the plasma is injected intravenously, and subsequent urine volume and electrolyte concentrations are noted. With blood from nonpregnant subjects, ninetyminute volumes are 17 to 42% of the levels at the time of plasma injection, but with the inhibiting factor proportions are 51 to 284%.

# Particle Size

#### AN INDEX TO READY DIGESTIBILITY



Sibly's STRAINED and HOMOGENIZED CARROTS

CARROTS MERELY STRAINED

THE ready digestibility of Libby's Strained AND Homogenized Baby Foods, and their early tolerability, are graphically shown as physical changes which Libby's exclusive process of homogenizing brings about.

For instance, in carrots that have only been strained, less than 30% of the food substance presents particles under 250 microns in size—more than 70% is composed of particles up to and over 840 microns in size. BUT when this substance undergoes Libby's homogenizing

process, there remain no particles over 250 microns in size; 87% are smaller than 150 microns.

Thus digestion is facilitated, and utilization of contained nutrients, such as iron, is enhanced. Since cellulose fibers are comminuted to ultrasmall size, Libby's Homogenized Baby Foods may be fed with safety as early as the fifth week of life and are well tolerated.\* Yet this feature carries no price penalty, for Libby's cost the mother no more than ordinary, merely strained, baby foods.
\*Reprints of clinical studies are appliable on request.

Libby, McNeill & Libby • Chicago 9, Illinois



HOMOGENIZED BABY FOODS



#### Neurology Spinal Fluid Cell Count in Air Encephalography

During air ventriculography with alternate withdrawal and replacement of cerebrospinal fluid, the cell count rises considerably. The height attained is in proportion to the original level. Mononuclear forms are largely responsible. Dr. Edwin R. Bickerstaff of the United Birmingham Hospitals, England, determined values at four stages of the procedure: at the start, after injection of 25 and 50 cc. of filtered air or oxygen, and at the end. Cells increased in 66 of 68 cases and rose within ten minutes in 50. Final levels were 30 to 92 cells per cubic millimeter in 22 instances. Averages for the last count ranged from 22 with 1 or no cells initially to 53 with 5 to 10 cells. Lancet 259:685-685, 1950.

#### Vital Statistics

#### U.S. Mortality

The lowest death rate in U.S. records, 9.6 per 1,000 population, is announced for 1950 by Dr. Louis I. Dublin, chief statistician of the Metropolitan Life Insurance Company. Mortality dropped 1% from 1949, with new low points for maternity, newborn, tuberculosis, and home accident levels and substantial gain in control of infection. Whooping cough was the only childhood disease in which fatalities failed to decline. Rises occurred for cancer. diabetes, chronic conditions of the heart, kidneys, and blood vessels, influenza, and accidents in automobiles and occupations.

#### Public Health Nursing Research

A five-year, million dollar research of nursing practice has been launched by the American Nurses' Association. From the outcome may be determined the amount and quality of nursing service required by the American people. Importance of the program is heightened by a crucial shortage of professional nurses in the national emergency. Studies will be conducted in hospitals under guidance of a committee representing medical, hospital, and women's organizations, as well as nursing fields. Chairman is the Rev. John J. Flanagan, S. J., Executive Director of the Catholic Hospital Association of the United States and Canada.

#### Epidemiology

#### Poliomyelitis Grants

Universities and research centers in sixteen states and Canada have been assigned approximately \$1,500,000 by the National Foundation for Infantile Paralysis for investigation and professional education in poliomyelitis. Approved projects will seek effective vaccine, a chemical agent to protect nerve cells from damage, a rapid diagnostic test, and antiserum that will increase individual resistance to paralysis. Funds will go to sixteen institutions for virus research, to nine for better methods of treatment, and to five for professional training. Since 1938, more than \$25,000,000 in March of Dimes collections has been spent for research and education, in addition to \$102,000,000 for patient care.


in acute follicular

tonsillitis . . .

"prompt clinical response"

with

Terramycin



CASE	DIAGNOSIS	CU	LTURE	DAILY	NUMBER OF	RESULT
	DIAGITOSIS	SOURCE	ORGANISM	DOSE GM.	DAYS TREATED	MESOET
29	Acute follicular tonsillitis	throat	Streptococcus pyogenes	4	3	Prompt clinical response. No fever after 24 hours of treatment

Case report taken from Herrell, W. E.; Heilman, F. R.; Wellman, W. E., and Bartholomew, L. A.: Proc. Staff Meet., Mayo Clin. 25:183 (Apr. 12) 1950

Dosage: On the basis of findings obtained in over 150 leading medical research centers, 2 Gm. daily by mouth in divided doses q. 6 h. is suggested for most acute infections. In severe infections, a high initial dose (1 Gm.) or higher daily doses (3 to 6 Gm.) should be used. Treatment should be continued for at least 48 hours after the temperature is normal and acute symptoms subside.

Supplied: 250 mg. capsules, bottles of 16 and 100;

100 mg. capsules, bottles of 25 and 100;

50 mg. capsules, bottles of 25 and 100,



Antibiotic Division

CHAS. PFIZER & CO., INC., Brooklyn 6, N. Y.

Muc



YDROPHILIC LUBRICOID

CHICAGO 11. ILLINOIS



# without oil

-presents methylcellulose as a gel, with magnesium hydroxide in less than laxative dosage to maintain hydration of the gel by osmosis.

#### Each tablespoonful contains:

METHYLCELLULOSE			 	*			6		,				0.3 см
MAGNESIUM HYDROXIDE.													0.6 GM.

#### The Turicum formula assures:

- lubricous bulk to encourage normal evacuation
- good distribution throughout the bowel
- no bloating
- no danger of impaction
- · no interference with utilization of oil-soluble vitamins
- no danger of lipid pneumonia
- no leakage

TURICUM IS AVAILABLE IN ONE PINT BOTTLES

#### Gastroenterology

#### Massive Gastric Hemorrhage

Gelfoam and thrombin swallowed at intervals after onset of severe gastroduodenal bleeding usually form a hemostatic clot. Dr. Meyer O. Cantor and associates of Detroit have treated 73 consecutive patients conservatively without a death, although the red cell count on admission was usually under 2,500,000 and shock often evident. If the source of hemorrhage is peptic ulcer, 2 tbs. of gelfoam mixed with 2 oz. of milk and cream is given orally every two hours and immediately followed by 250 units of thrombin in 50 cc. of a water solution. Amphojel is administered at the rate of 1 tbs. every three hours. Additional measures include blood replacement in amounts up to 10,000 cc. in twentyfour hours, Meulengracht's diet, antacid therapy, and sedation. For bleeding esophageal varices, gelfoam powder is given in the dry state and followed only by thrombin solution. Operation should be done promptly after exsanguinating hemorrhage or failure of blood transfusion to improve circulation within twenty-four hours.

Am. J. Surg. 80:883-887, 1950.

#### Honors

#### Public Health Award

Dr. Rolla E. Dyer of Emory University, Atlanta, former director of the National Institutes of Health, has won the Sedgwick Memorial Medal presented annually by the American Public Health Association for distinguished service. He did outstanding work on Rocky Mountain spotted fever, typhus, and Q fever.

#### Doctor to Doctor

Think of a gag that fits the illustration. For every issue a new gag is published and the author is sent \$5. The Feb. 15 winner is J. F. Fulp, M.D.

Stoneville, N. C.
Mail your caption to
The Cartoon Editor
Caption Contest
No. 2

Modern Medicine 84 South 10th St. Minneapolis 3, Minn.

176



"So what if they got your 80%? You should smile. Mr. Truman says next year it's really going to hurt."

# 3

# Useful Cardiac Drugs

#### • Thesodate - Brewer IN ANGINA PECTORIS

(Theobromine Sodium Acetate 71/2 gr. enteric coated)

Thesodate has been proven effective in increasing the capacity for work in individuals suffering from coronary artery disease. One Thesodate tablet four times a day (after meals and at bedtime) helps to maintain improved heart action and increased coronary artery circulation.

#### € Enkide - Brewer IN LUETIC HEART DISEASE

(Potassium Iodide one gram or half gram enteric coated)

Enkide is useful as an adjuvant in tertiary syphilis and wherever potassium iodide therapy is indicated. Enkide insures accuracy of dosage, absence of gastric irritation and convenience of administration. Patients are more apt to follow prescription directions because of these advantages.

#### ♠ Amchlor - Brewer IN CARDIAC EDEMA

(Ammonium Chloride one gram enteric coated)

Amchlor cuts in half the number of tablets each patient takes when large amounts of ammonium chloride are prescribed. This convenience to the patient helps to insure full and complete use of the entire amount prescribed. Amchlor is useful in cardiac edema, hypertension, dysmenorrhea, Meniere's Syndrome, etc.

For samples - just send your Is blank marked MM-2.



BREWER & COMPANY, INC. WORCESTER 8, MASSACHUSETTS U.S.A.

#### Chemotherapy

#### Drugs for Peritonitis

Death from widespread polymicrobial peritonitis in dogs may be averted by intramuscular injection of penicillin in beeswax and oil at intervals of twelve hours. Mortality is lower than when penicillin is combined with streptomycin or sulfonamides, alone or together, or when only streptomycin is used, report Dr. H. A. Zintel and associates of the University of Pennsylvania, Philadelphia. The various types of chemotherapy were employed for dogs after contamination of the peritoneum with contents of the incised appendix, which was ligated and left in the abdomen. The survival rate varied from 66% with penicillin to 27% with streptomycin and was 3.7% with no treatment. Recovery seemed unrelated to bacterial types or strains noted before or after therapy or to antibiotic sensitivities of organisms in the first cultures.

Surg., Gynec. & Obst. 91:742-750, 1950.

#### Radiology

#### Radiation Death Prevention

Antibiotics may lower mortality of the radiation syndrome to a third or a fifth of expected rates, after exposure commonly lethal in 70 to 100% of cases. Sickness is also reduced and, with fatal injury, survival time is prolonged, although actual damage to tissues is not influenced. On noting that rats killed by injection of radioactive phosphorus had both gram-negative and gram-positive bacteria in the lesions, Drs. Simon Koletsky and James H.

Christie of Western Reserve University, Cleveland, used penicillin and streptomycin in combination for protection. Benefits were similar whether intramuscular doses were given for three to ten days before injection of Pa2 in different amounts or immediately afterward in courses not exceeding three weeks. Mortality was lowered in 8 of 9 groups and in 4 was greatly reduced, for example, from 67 to 14% and from 100 to 33%. Weight loss, diarrhea, and bleeding were also distinctly less in treated animals.

Proc. Soc. Exper. Biol. & Med. 75:363-366, 1950.

#### Surgery

#### Prevention of Adhesions

When postoperative intestinal adhesions have been cut apart, regrowth is far less likely if motility of the bowel is maintained by early feeding and injections of Prostigmine. The regimen is particularly suitable if surgery is extensive but does not include intestinal resection. In preliminary trial on dogs, Dr. C. A. Schiff and associates of Michael Reese Hospital, Chicago, stimulated tissue formation by distributing talcum powder over the small bowel and, six weeks later, severed fibrous bands. For the first two days after operation some animals received subcutaneous injections of Prostigmine methylsulfate every four hours and were encouraged to eat and drink, while others had atropine sulfate and nothing by mouth. About 33% of adhesions re-formed in the first group and 92% in the second. Surgery 28:977-982, 1950.

# How mild can a cigarette be?

• Every day, more and more smokers - and among them many, many doctors - are discovering for themselves just how mild a cigarette can be. They're making their own 30-Day Camel Mildness Tests-smoking Camels regularly for 30 days.

It's a sensible cigarette test. As a doctor, you know there can be no valid conclusion drawn from a one puff comparison from a trick test that calls for hasty decisions. The Camel 30-Day Test asks you to make a day after day, pack after pack comparison.

If you are not already a Camel smoker, why not try this test? Judge Camel mildness and the rich, full flavor of Camel's choice tobaccos in your own "T-Zone"the real proving ground for a cigarette.

Make your own 30-Day Camel Mildness Test in your own "T-Zone"-That's T for Throat, T for Taste. See if you don't change to Camels for keeps!



More People omoke Camela
THAN ANY OTHER CIGARETTE!

#### Nutrition

#### Antianemic Factor

Folinic acid, commonly termed the citrovorum factor, when given in adequate amounts, promptly induces blood regeneration and relieves symptoms of pernicious anemia, nutritional macrocytic anemia, and tropical sprue in relapse. The superiority of large doses was demonstrated in 10 cases by Dr. Tom D. Spies and associates of Northwestern University, Chicago, Hillman Hospital, Birmingham, and General Calixto Hospital, Havana. Garcia Daily amounts of 200 units administered for ten days had insufficient effect in 2 instances and none in 4. The remaining 4 patients received intramuscular injections of 20,000,000 units per day for the same period, with excellent response in hematopoiesis and well-being. A dose of 10,000,000 units equals 1 mg. of pteroylglutamic acid.

South. M. J. 43:1076-1082, 1950.



#### Vital Statistics

#### U.S. Mortality

In every age group, the death rate for males is higher than that for females, and the latter is falling more rapidly, according to figures of the U.S. Public Health Service. Between 1940 and 1949, mortality of women and girls dropped 13% and of men and boys 7%, to 8.3 and 11.2 per 1,000 respectively. The largest relative decreases took place in younger groups. For example, the rate for 1 to 14 years declined 40%, and for 65 to 74 years 10%. The greatest sex difference was in the 15- to 24year division, where the excess for males was 89%, more than 3 times the difference for 1940.

#### Publications

#### Journal Gets New Editor

The American Journal of the Medical Sciences is now under the editorial direction of Dr. Richard A. Kern of Temple University, Philadelphia. He is assisted by Drs. Thomas M. Durant and Chris J. D. Zarafonetis. Dr. E. B. Krumbhaar of the University of Pennsylvania, who served twenty-five years as chief editor, retired at the year's end. Lea and Febiger are publishers of the journal.

#### Honors

#### Chemical Society Award

Dr. Roger J. Williams of the University of Texas, Austin, discoverer of the growth-promoting vitamin pantothenic acid, and a pioneer in folic acid therapy of pernicious anemia, has received the American Chemical Society Southwest award.

# so comfortable

Both physically and psychologically, TAMPAX tampons are amazingly comfortable intravaginal menstrual guards. They cannot induce odor, perineal irritation or infection via rectum. And, with the individualization and convenience of protection provided by the three absorbencies (Regular, Super, Junior), their use is said to tend to make women "forget they are menstruating."\* These dainty cotton tampons are also thoroughly safe and adequate.

\*West. J. Surg., Obstet. & Gynec., 51:50, 1943; J.A.M.A., 128:490, 1945.

TAMPAX INCORPORATED PALMER, MASS.

#### TAMPAX

the internal menstrual guard of choice

Your request will bring professional samples promptly.



MM-15-21

#### Current Books & Pamphlets

This catalogue is compiled from all available sources, American and foreign, to insure a complete listing of the month's releases.

#### Medicine

THE SCOURGE OF RHEUMATISM, REPORT OF A CONFERENCE HELD BY THE BRITISH RHEUMATIC ASSOCIATION IN SEPTEMBER, 1949. 85 pp. William Heinemann Medical Books, London. 5s.

VIRUSES 1950 edited by M. Delbrück. 147 pp., ill. California Institute of Tech-

nology, Pasadena. \$2.50

DIE ANSTECKENDEN KRANKHEITEN: IHRE EPIDEMIOLOGIE, BEKÄMPFUNG UND SPE-ZIFISCHE THERAPIE edited by Max Gundel. 4th ed. 1,056 pp., ill. Georg Thieme, Stuttgart. 66 DM.

PROGRESS VOLUME: MODERN DEVELOPMENTS IN THERAPEUTICS AND METHODS OF TREATMENT by Harold Thomas Hyman. 867 pp. W. B. Saunders Co.,

Philadelphia. \$10

PHYSICAL EXAMINATION IN HEALTH AND DISEASE by Rudolph H. Kampmeier. 821 pp., ill. F. A. Davis Co., Philadelphia. \$8

SEROLOGY WITH LIPID ANTIGEN: WITH SPE-CIAL REFERENCE TO KAHN AND UNIVER-SAL REACTIONS by Reuben L. Kahn. 327 pp., ill. Williams & Wilkins Co., Baltimore. \$6

LE RÉTRÉCISSEMENT MITRAL: ÉTUDES ANAT-OMIQUES, CLINIQUES ET THÉRAPEUTIQUES by R. Lutembacher. 304 pp., ill. Mas-

son & Co., Paris, 1,400 fr.

#### Neurosurgery

STUDIES IN LOBOTOMY edited by Milton Greenblatt, Robert Arnot and Harry C. Solomon. 495 pp., ill. Grune & Stratton, New York City. \$10

CRANIOPLASTY by David L. Reeves. 119 pp., ill. Charles C Thomas, Spring-

field, III. \$3

#### Physiology

PHYSIOLOGY AND ANATOMY by Esther M. Greisheimer. 6th ed. 841 pp., ill. J. B. Lippincott Co., Philadelphia. \$4

COMPARATIVE ANIMAL PHYSIOLOGY David W. Bishop et al.; edited by C. Ladd Prosser. 888 pp., ill. W. B. Saunders Co., Philadelphia. \$12.50

#### Embryology

CHEMICAL EMBRYOLOGY by Jean Brachet. 533 pp., ill. Interscience Publishers, New York City. \$8

HUMAN EMBRYOLOGY FOR MEDICAL STU-DENTS by S. R. Nair. 398 pp., ill. The Popular Book Depot, Bombay. 425. OF 28 TS.

#### Biochemistry

BACTERIAL POLYSACCHARIDES: THEIR CHEMI-CAL AND IMMUNOLOGICAL ASPECTS by Martin Burger. 273 pp. Charles C Thomas, Springfield, Ill. \$6

CELL PHYSIOLOGY AND PHARMACOLOGY by James F. Danielli. 156 pp., ill. Elsevier Publishing Co., London. 24s.

CHEMISTRY AND BIOLOGY OF PROTEINS by Felix Haurowitz. 374 pp., ill. Academic Press, New York City. \$5.50

A TEXTBOOK OF BIOCHEMISTRY by Philip Henry Mitchell. 2d ed. 695 pp., ill. McGraw-Hill Book Co., New City. \$6

MEDIZINISCHE CHEMIE: EIN LERN-HILFSBUCH FÜR MEDIZINER by Philip Fritz Vering. 493 pp., ill. Wilhelm Maudrich, Vienna. 150 Sch.

THE BIOCHEMISTRY OF B VITAMINS by Roger John Williams et al. 741 pp., ill. Reinhold Publishing Corp., New

York City. \$10

# Penicillin-Triple Sulfonamide Mixture in Palatable Liquid Form...



For additive or possible synergistic action . . .

# Triple Sulfonamides

—in pleasant-tasting, convenient liquid form, readily acceptable to infants, children, adults.

White's Dram-cillin with Triple Sulfonamides contains, per teaspoonful (5 cc.): 100,000 Units of Penicillin G Potassium and 0.5 grams of triple sulfonamide mixture: equal parts of

Sulfadiazine Sulfamerazine Sulfacetimide (buffered)

Greater Safety—Oral penicillin appears less likely to produce hypersensitivity reactions. Combined sulfonamides permit smaller, safer dosage of each<sup>2</sup>—offer less likelihood of crystalluria.

#### Supplied: In 60 cc. bottles

ALSO AVAILABLE: Potent Oral Penicillin for All Age Groups

White's Drop-cillin: 50,000 units\* in a dropperful.
9 cc. bottle contains 600,000 units.

White's Dram-cillin: 100,000 units\* in a teaspoonful. 60 cc. bottle contains 1,200,000 units.

Both with a delicious vanilla flavor.

\*buffered penicillin G potassium

#### WHITE LABORATORIES, INC., Pharmaceutical Manufacturers, Newark 7, N. J.

- 1. Keefer, C. S.: Amer. J. Med. 7:216 (1948).
- 2. Kolmer, J. A.: Amer. J. Med. Sc. 215:136 (1948).

#### Anatomy

AN ATLAS OF HUMAN ANATOMY by Barry Joseph Anson. 518 pp., ill. W. B. Saunders Co., Philadelphia. \$11.50

THE ESOPHAGUS AND PHARYNX IN ACTION: A STUDY OF STRUCTURE IN RELATION TO FUNCTION by William Lerche. 222 pp., ill. Charles C Thomas, Springfield, Ill. \$5.50

MANUAL FOR COMPARATIVE ANATOMY by Leonard P. Sayles. Rev. ed. 214 pp., ill. Macmillan Co., New York City. \$3

FASCHENBUCH DER ANATOMIE, VOL. 1, by Hermann Voss and Robert Herrlinger. 3d ed. 344 pp., ill. Gustav Fischer, Jena. 10 DM.

#### Military Medicine

UNDER THE DIRECTION OF THE LOS ALAMOS SCIENTIFIC LABORATORY edited by J. O. Hirschfelder et al. 456 pp., ill. McGraw-Hill Book Co., New York City. \$3

GERMAN AVIATION MEDICINE, WORLD WAR II. PREPARED UNDER THE AUSPICES OF THE SURGEON-GENERAL, U.S. AIR FORCE. 2 vols., 1,302 pp., ill. U.S. Government Printing Office, Washington, D.C. \$8,50



"Ready for the next kid, Doctor?"

#### Gynecology & Obstetrics

MODERN TRENDS IN OBSTETRICS AND GYNAE-COLOGY edited by Robert Kenneth Bowes. 778 pp., ill. Butterworth & Co., London. 60s.; Harper & Brothers, New York City. \$12

PERMANENT CONCEPTION CONTROL by Robert L. Dickinson and Clarence J. Gamble. 39 pp., ill. Planned Parenthood Federation of America, New York City. Apply

YOU'LL LIVE THROUGH IT: FACTS ABOUT THE MENOPAUSE by Miriam Lincoln. 182 pp., ill. Harper & Brothers, New York City. \$2.50

FERTILITY IN MARRIAGE: A GUIDE FOR THE CHILDLESS by Louis Portnoy and Jules Saltman. 250 pp., ill. Farrar, Strauss & Co., New York City. \$3

J. P. Pundel. 350 pp., ill. Masson & Co., Paris. 2,000 fr.

INTRODUCTION TO MOTHERHOOD by Grantley D. Read. 92 pp., ill. William Heinemann Medical Books, London.

VOUR PREGNANCY by Anna Roosevelt and Leo Doyle. 178 pp. Henry Holt & Co., New York City. \$2.50

UNDERSTANDING NATURAL CHILDBIRTH: A BOOK FOR THE EXPECTANT MOTHER by Herbert Thoms and Laurence G. Roth. 112 pp., ill. McGraw-Hill Book Co., New York City. \$3.50

CHIRURGISCH-GYNAKOLOGISCHE GRENZFRA-GEN: PHYSIOLOGIE UND PATHOLOGIE DER WEIBLICHEN GESCHLECHTSORGANE IN BEZIEHUNG ZUR CHIRURGIE by Herbert Uebermuch. 168 pp., ill. Georg Thieme, Leipzig. 9 M.

#### Sex

- SEXUAL BEHAVIOR IN SOCIETY by Alex Comfort, 157 pp. Viking Press, New York City. \$2.75
- WHEN YOUNG PEOPLE ASK ABOUT SEX by
  Dora Hudson Klemer. 62 pp., ill.
  Association Press, New York City. 75¢
  MARRIAGE IS WHAT YOU MAKE IT by
- MARRIAGE IS WHAT YOU MAKE IT by Paul Popenoe. 221 pp. Macmillan Co., New York City. \$3



#### **Normal Functions Regained**

The inability to perform such simple everyday tasks as combing hair, pulling up a girdle, or tying a shoelace may rob arthritics of self-reliance and independence.

DARTHRONOL, through the proven antiarthritic effects of massive dosage Vitamin D, and the important nutritional benefits of 8 vitamins, is instrumental in abolishing pain, diminishing soft tissue swelling, restoring normal function and promoting an improved sense of well-being in an ever-increasing number of disabled arthritics.

# Darthronol

FOR THE




#### BACH CARSHIE CONTAINS

Vitomin	D (Irradiated Ergosteral) 50,000 USP unit
Vitamin	A (Fish Liver Oil) 5,000 USP unit
Vitamin	C (Ascorbic Acid)
Vitamin	B, (Thiamine Hydrochloride)3 mg
Vitamin	8: (Riboflavin)
Vitamin	B. (Pyridoxine Hydrochloride)0.3 mg
Niecine	mide
	Pantothenate
Mixed T	ocopherols 4 mg
	ent to 3 mg. of Synthetic Alpha Tocopheral



J. B. ROERIG AND COMPANY . SIS CASE SHORE DR. CRICAGO IS ICLINOIS



#### the modern therapy in VAGINITIS

50% of all women patients exhibit vaginal discharge, and pH is the index.

Since acidity is the most important therapeutic factor in vaginitis, DOMOGYN, with an optimum pH of 4.2 quickly and effectively controls the condition and reestablishes optimum acidity, and a normal vaginal flora.

One teaspoonful of bulk DOMOGYN, or the contents of a DOMOGYN Packette in 2 quarts of warm water as a soothing therapeutic douche.

Available at all drug stores



DOME CHEMICALS, INC. 100 West Bath St. New York 73 M T

#### WHEN A HEARING

Is Necessary



-you can rely on PARAVOX to supply ample volume, clearness of tone, and other desirable qualities, all backed by a widespread service plan. The PARA-VOX Tiny-MYTE Hearing Aid shown above is being worn by thousands with satisfaction.

PARAVOX Hearing Aids are widely known in the medical profession and were exhibited at the San Francisco Annual Session of the AMA, the Annual Meetings of the American Academy of Ophthalmology and Oto-

laryngology, the Medical Society of Penn., and the Mississippi Valley Medical Society.



2056 E. 4th St. Cleveland 15, Ohio

#### Ophthalmology

PRINCIPLES OF OPHTHALMOLOGY by Thomson Henderson. 224 pp., ill. William Heinemann Medical Books, London.

IN MY MIND'S EYE by Frederick Marion. 315 pp. E. P. Dutton & Co., New York City. \$3.75

RESEARCHES IN BINOCULAR VISION by Kenneth N. Ogle. 345 pp., ill. W. B. Saunders Co., Philadelphia. \$7.50

#### Essays

APHORISMS: FROM THE BEDSIDE TEACHINGS AND WRITINGS OF SIR WILLIAM OSLER collected by Robert Bennett Bean. edited by William Bennett Bean. 159 pp. Henry Schuman, New York City. \$2.50

DOCTOR COME QUICKLY by Frank J. Clancy. 248 pp. Superior Publishing Co., Seattle. \$2.95

HUMANISM IN AN AGE OF SCIENCE: BEING A LUDWIG MOND LECTURE DELIVERED AT THE MANCHESTER SCHOOL OF MEDICINE ON 6 OCTOBER 1949 by John F. Fulton. 26 pp. Henry Schuman, New York City. 75¢

ESSAYS IN THE HISTORY OF MEDICINE by George E. Gask. 209 pp., ill. Butterworth & Co., London. 30s.

THE DOCTOR TAKES A FARM by Jeff Minckler. 90 pp. Dorrance & Co., Philadelphia. \$2.50

MEDICINE AND MANKIND by Arnold Sorsby. 2d ed. 232 pp., ill. Staples Press, London. 12s. 6d.

A DOCTOR DIGRESSES by K. Walker. 207 pp. Jonathan Cape, London. 10s. 6d.



# When you recommend steam therapy consider Vicks VapoRub as the medicament

You can increase the benefits your patients derive from steam inhalation by suggesting Vicks VapoRub as the medicament.

Its well-balanced formula contains not one but seven volatilizing ingredients, including menthol, thymol, camphor and oil of eucalyptus—all helpful in soothing the irritated mucosa of the respiratory tract, as well as in combatting dryness.

So consider Vicks VapoRub
when your patients require
steam therapy, whether you
recommend a vaporizer or
some other method.
In practically every
home, Vicks VapoRub
is already on hand
for instant use.

### for your patients

We will be happy to send you a generous supply of distribution samples. Why not use this handy coupon?



VICK CHEMICAL COMPANY Department N, Box 1813 Greensboro, North Carolina

Please send me, without obligation, a supply of distribution samples of Vicks VapoRub.

Name\_\_\_\_

Street

City\_\_\_\_State\_\_\_\_

## ANPLEX Chimedie

In all types of anemia, ANPLEX (Chimedic) assures rapid hemoglobic regeneration and effective stimulation of reticulocytosis through the potent antianemic action of: Vitamin Big. U.S.P. (Crystalline)—Folic Acid—Crude Liver.

ANPLEX (Chimedic) is safe and nonirritating on injection.

#### LITERATURE ON REQUEST

#### CHICAGO PHARMACAL COMPANY

5547 N. Ravenswood Ave., Chicago 40, III.



Cases: Excellent results:

Each Flavette contains 1/20
gr benzocaine with flavoring
phetamine, or other potent
clinical reprints
N Y St J Med May '47—"N Carolina M J July '50
AMHERST RESEARCH DIVISION
Capitol Station, Albany, N. Y.

#### For Patients Suffering From



Prescribe Dr. Scholl's Arch Supports in cases requiring mechanical relief from Foot Arch Trouble of any kind. The patient will be properly fitted and the Supports adjusted at no extra cost as the condition of the foot improves. This nation-wide Service is available at many leading Shoe and Dept. Stores and at Dr. Scholl's Foot Comfort® Shops in principal cities.

Dr Scholl's SUPPORTS

#### PATIENTS

#### . . I Have Met

The editors will pay \$1 for each story published. No contributions will be returned. Send your experiences to the Patients I Have Met Editor, MODERN MEDICINE. 84 South Tenth St., Minneapolis 3, Minn.

#### That's What the Man Said

When I was in medical school we used to have fun spoofing one of the janitors in the lab building. We had been running some urine tests when the janitor came in. I poured the contents of two or three tubes into a beaker and handed it to John.

ed it to John.
"Here," I said, "drink this. If it kills you we'll hold an autopsy."

"Over my dead body, you will," retorted the janitor.

"Yep, that's what he said," quipped my partner.—o.p.a.

#### Sounded Different

One of our patients is a Miss Laub. I am sure our receptionist did not mean to be derogatory when she announced, "Miss Slob is waiting!"—A.S.L.



"She went to pieces when the doctor told her there was nothing the matter with her."



# Thinner than your skin —and stronger

WHEN a pair of B. F. Goodrich surgeons' gloves are stretched over your hands, the gloves are actually thinner than the skin they cover.

B. F. Goodrich surgeons' gloves, made by the patented Anode process, keep your touch almost as sensitive as it would be without gloves. When operating, you can forget any worry about your gloves because they are made in a way that completely eliminates weak spots. Between

the fingers, at the fingertips and at the wrist, normally the weakest spots, these single-layer gloves stay uniformly strong.

The extra long, restricted wrists, full backs and tapered fingers all add up to your comfort when operating or examining. A full range of accurate sizes assures you the exact fit you need.

These gloves come in three types: surgeons' gloves for operating, short wrist examining gloves and "Special Purpose" gloves which were created for those who develop an allergic dermatitis when wearing ordinary rubber gloves.

Order B. F. Goodrich gloves from your hospital or surgical supply dealer. The B. F. Goodrich Company, Sundries Division, Akron, Ohio.

## B.F. Goodrich Surgeons' Gloves

#### now in a single preparation:

More and more physicians are turning to combined penicillin-sulfonamide therapy; for it has now been demonstrated that in many instances this combination is more effective than—and offers many advantages over—the newer antibiotics.

'Eskacillin-Sulfas' is leading the trend to this new and exciting combination therapy because it is presented to you in the most logical, easily-given and readily-accepted form: an exceptionally palatable fluid.

'Eskacillin-Sulfas' is for the treatment of infections caused by organisms sensitive to the action of penicillin or the sulfonamides.

Among its many indications are:

Bronchitis	Pneumonia	Sinusitis				
Tonsillitis	Otitis Media	Gonorrhea				

You will also find 'Eskacillin-Sulfas' especially useful in urinary tract infections; bacillary dysentery; meningococcic infections; prophylaxis in streptococcal infections and rheumatic fever; as adjunctive therapy in pneumococcal meningitis and H. influenzae meningitis after the acute phase has subsided; prophylaxis in colonic surgery, surgical wounds and burns; and in certain resistant staphylococcal infections.

Due to its extremely wide antibacterial spectrum and the minimum chance of the development of resistant strains, many physicians find 'Eskacillin-Sulfas' the ideal therapy when immediate medication is necessary and lack of either time or laboratory facilities prevents exact determination of the specific organism involved.

## 'ESKACILLIN-SULFAS'

the original presentation of <u>penicillin</u> and the <u>sulfonamides</u> in fluid form Available in 2 fl. oz. bottles

Smith, Kline & French Laboratories, Philadelphia

#### why 'Eskacillin-Sulfas' is in fluid form:

We concluded—after weighing every possibility—that 'Eskacillin-Sulfas' should be a *fluid* for the following reasons:

- Tablets are impractical because the bulky nature of the sulfonamides makes it virtually impossible to manufacture a small, easily-swallowed tablet.
- 2. The fluid form assures more rapid attainment of therapeutic blood levels.
- A palatable fluid is the ideal form for children and for the many adults who balk at, or have difficulty with, tablet medication.

'Eskacillin-Sulfas' is so pleasant tasting your patients will find it easy to take whatever amount you prescribe—and, because of its fluid form, you will find it easy to adjust the dosage.







Codeine and hyoscyamus plus ammonium hypophosphite, white pine and tolu in a glycerin base provide sedation of the cough reflex — liquefy mucus. General dosage: Adults 1 to 2 teaspoonfuls every 2 to 3 hrs. Children in proportion. Literature available to physicians.

MARTIN H. SMITH COMPANY

#### Stitch in Time

An angry young father burst into my office and brandished a bruised fist in my face.

"Come, come," I said, "take it easy. What seems to be the matter?"

"Matter?" he echoed. "If you'd been around last night you would have found out. I would have hit you instead of the door. What did you expose my wife for?"

It took a little doing but I finally convinced him that it was my regular practice to make such an examination after childbirth when stitches have to be removed.—R.G.F.

"Yes, we'll admit your T & A case and schedule the operation for tomorrow morning, but our board man will do it for you."—M.P.C.



"The tests show that your son is allergic to soap and water."

#### Let's Change the Subject

The patient had come to a psychologist for help because she blushed so easily.

"Why, when I just sit down and think, I blush," she said. "What can I do to stop it?"

"Probably the best way," said the psychologist, "would be to think of something different."—C.R.A.

# New...

# New LOW PRICES New TABLET POTENCIES

A Substantial Saving to Your Patients Receiving....

# **VERILOID**\*

#### IN THE TREATMENT OF HYPERTENSION

The excellent acceptance accorded Veriloid by the profession, and the construction of expanded manufacturing facilities have made possible a substantial reduction in the price of this unusual hypotensive agent.

#### TWO NEW POTENCIES

In addition to the formerly available 1 mg. tablet, Veriloid is now also available in 2 mg. and 3 mg. tablets. Based on former prices, the 1 mg. tablet represents a saving of  $16\frac{2}{3}\%$ , the 2 mg. tablet a saving of 25%, and the 3 mg. tablet a saving of  $33\frac{1}{3}\%$ .


After the optimal dose has been determined, the prescribing of the largest tablet size will result in the greatest saving. Literature describing the action, uses, and administration of Veriloid is available on request.

\*Trade Mark of Riker Laboratories, Inc.

RIKER LABORATORIES, INC. 8480 BEVERLY BLVD., LOS ANGELES 48, CALIF.



Veriloid, a purified biologically standardized extract of Veratrum viride, is now supplied in 1.0 mg. 2.0 mg., and 3.0 mg. scored tablets in bottles of 100, 500, and 1000. Available on prescription only at all pharmacies.



# PASTER RUB GIVES FASTER RELIEF IN RHEUMATIC ACHES-PAINS

**Lumbage and Neuritis Discomfort** 

Musterole is an excellent analgesic, decongestive, and counter-irritant for relieving muscular aches, pains, sorenees and stiffness—for helping to break up topical congestion. It has all the advantages of a mustard plaster yet eliminates the fuss and bother of making one, and is far more comfortable for the patient. In 3 Strengths: Children's Mild. Also Regular and Extra Strong for adults.



#### INDEX TO ADVERTISERS

Airkem, Inc. Alden, John, Tobacco Company Alkalol Co, The American Cystoscope Makers, Inc. American Ferment Company, Inc. Amers Company, Inc. Amers Company, Inc. Amers Mekenna & Harrison Limited Baby Bathinette Corporation Barnes, A. C., Company Bauer & Black Becton, Dickinson & Co. Birtcher Corporation, The Brewer & Company, Inc. Carnrick, G. W. Company Chicago Pharmacai Co. Ciba Pharmaceutical Products, Inc. Crockes Laboratories Inc. Cutter Laboratories. Davol Rubber Company	135
Alkalol Co. The	186
American Cystoscope Makers, Inc	139
American Ferment Company, Inc	23
Amherst Research Division	188
Armstrong Cork Co	. 22
Ayerst, McKenna & Harrison Limited	37
Barnes A C Company	192
Bauer & Black	113
Becton, Dickinson & Co	143
Birtcher Corporation, The	163
Carnrick, G. W., Company	58
Chicago Pharmacal Co	188
Ciba Pharmaceutical Products, Inc4th	Cover 33
Crookes Laboratories, Inc.	32
Cutter Laboratories	105
Davol Rubber Company	151
De Vilbias Company. The	147
Dietene Company, The	47
Dome Chemicals, Inc.	186
Fleet. C. B. Co. Inc.	55
Plorida Citrus Commission	159
Geigy Company, Inc	115
Goodrich B. F. Company. The	189
Hanovia Chemical & Manufacturing Co	46
Harrower Laboratory, Inc., The	167
Hoffmann-La Roche Inc	50-51
International Pharmaceutical Corp	62
Irwin, Neisler & Co	129
Kinney & Company	141
Knox Gelatine	133
Kremers-Urban Company	155
Commercial Solvents Corporation Crookes Laboratories, Inc. Cutter Laboratories, Inc. Cutter Laboratories. Davol Rubber Company Denver Chemical Mfg. Co. Inc. De Vibibis Company, The Dietene Company, The Dietene Company, The Dome Chemicals, Inc. Ethicon Suture Laboratories, Inc. Fleet, C. B., Co., Inc., Florida Citrus Commission Geigy Company, Inc., General Electric X-Ray Corp., Goodrich, B. F., Company, The Hanovia Chemical & Manufacturing Co. Harrower Laboratory, Inc., Heinz, H. J., Co. Hoffmann-La Roche, Inc. International Pharmaceutical Corp. Irwin, Neisier & Co. Ivers-Lee Company Knox Gelatine, Kremers-Urban Company Lakeside Laboratories, Inc. Leeming Thos. & Co. Inc. Leeming Thos.	161
Libby, McNeill & Libby	171
Aremers-Uroan Company Lakeside Laboratories, Inc. Leeming, Thos., & Co., Inc Libby, McNeill & Libby Lilly, Eli & Company Mallinckrodt Chemical Works Matthe Laboratories, Inc. McNeil Laboratories, Inc.  McNeil Laboratories, Inc.  55	8
Mallinckrodt Chemical Works	52-153
McNeil Laboratories, Inc.	14-15
Mead Johnson & Company	9, 125
Merck & Co., Inc.	Cover
Musterole Co	
	194
National Drug Company, The	117
National Drug Company, The	43
National Drug Company, The	117
McNeil Laboratories, Inc.  Mead Johnson & Company 5  Merck & Co., Inc.  Merrell, The Wm. S. Company 2nd  Musterole Co.  National Drug Company, The.  National Electric Instrument Co. Inc.  Nepera Chemical Co., Inc.  Numotizine, Inc.  Owens-Corning Piberglas Corporation	
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc.	
National Drug Company, The. National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company Patch, The E. L. Company	
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company Patch, The E. L., Company Pelton & Crane Company, The	
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company, Patch, The E. L., Company, Pelton & Crane Company, The Pet Milk Company, Pet Milk	
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company Patch, The E. L., Company, Pelton & Crane Company, The Pet Milk Company, Pfizer, Chas. & Co., Inc. 22 Rand Pharmaceutical Co. Inc.	
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company Patch, The E. L., Company, Pelton & Crane Company, The Pet Milk Company Pizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Reynolds, R. J., Tobacco Company	117 43 41 119 165 186 13 21 127 173 28 179
National Drug Company, The National Ricctric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company Patch, The E. L. Company Pelton & Crane Company, The Pet Milk Company Pizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Reynolds, R. J., Tobacco Company Riker Laboratories, Inc.	117 43 41 119 165 186 13 
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Parke, Davis & Company, Patch, The E. L. Company, Pelton & Crane Company, The Pet Milk Company, The Pet Milk Company, Co., Inc. Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Robrins, J. B. & Company, 16	117 43 41 119 165 186 13 
National Drug Company, The National Electric Instrument Co., Inc. Nepera Chemical Co., Inc. Numotizine, Inc. Numotizine, Inc. Owens-Corning Piberglas Corporation Paravox, Inc. Paravox, Inc. Parke, Davis & Company Patch, The E. L., Company, Pelton & Crane Company, Pelton & Crane Company, Piter, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company, Stepany, Stepa	117 43 41 119 165 186 13 3 121 121 13 3 121 193 188 169 188 189 189 189 189 189 189 189 189 18
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L., Company, Pelton & Crane Company, The Pet Milk Company. Pfizer, Chas., & Co., Inc. Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. Robins, A. H. Company, Inc. Royal Metal Mfg. Company Sanborn Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 18-169 185 29 24
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 13 3 121 121 13 3 121 193 188 169 188 189 189 189 189 189 189 189 189 18
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 18-169 185 29 24 36 61 196
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 13 121 127 7, 173 8 179 193 18-169 185 29 24 66 61
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 18-169 185 29 24 36 61 196
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 133 3 121 127 5, 173 38 169 193 18-169 29 24 61 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 133 121 127 5, 173 38 179 193 38-169 185 29 24 36 61 196 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company, Pelton & Crane Company, The Pelton & Crane Company, The Pel Milk Company, Pfizer, Chas. & Co., Inc. Rand Pharmaceutical Co., Inc. Royal Metal Mfg. Company, Sanborn Company, Sanborn Company, Sandoz Pharmaceuticals Schering Corporation Schmid, Julius, Inc. Scholl Mfg. Co., Inc., The Seeck & Kade, Inc. Skiar, J., Manufacturing Co. Smith, Kline & French Laboratorica T., 17, 19, 20, 145, 157, 19 Smith Martin H. Company	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192
Owens-Corning Piberglas Corporation Parawox, Inc. Parke, Davis & Company Patch, The E. L. Company. Pelton & Crane Company, The Pet Milk Company. Pifzer, Chas. & Co., Inc. 2: Rand Pharmaceutical Co., Inc. 2: Rand Pharmaceutical Co., Inc. 3: Reynolds, R. J., Tobacco Company, Riker Laboratories, Inc. Robins, A. H. Company, Inc. 30-31, 16 Roerig, J. B., & Company. Sanborn Company, Sanborn Company. Sanborn Company.	117 43 41 119 165 186 13 3 121 127 5, 173 38 179 193 8-169 24 24 24 26 61 188 42 192

#### THE TABLET SHE CAN TOLERATE



ULVICAL "Ulmer" combines all the prenatal patient's requirements into one easy-to-take, well-tolerated, economical tablet. As a dietary supplement during pregnancy and lactation, it supplies sufficient iron to correct or prevent any tendency toward anemia, minimum daily requirements of calcium and all the essential vitamins.

FOR	RMULA
Vitamin A	Vitamin E (A-tocopherol)
Vitamin D 200 U.S.P. Units	Calcium Pyrophosphate 7.5 grains
Thiamine HCI	represents Calcium 150 mg.
Riboflavin	Phosphorus 120 mg.
Ascorbic Acid	Ferrous Sulfate U.S.P

To eliminate the possibility of intolerance, the ascorbic acid and thiamine hydrochloride are released and utilized in the stomach while the other factors are not released until they enter the small intestine. Endorsed and prescribed by leading obstetricians, ULVICAL "Ulmer" provides assured results in the recommended dosage of 2 to 6 tablets per day.



State

City

W

IMMOBILIZES
SPERM IN THE
FASTEST TIME
RECOGNIZED
FOR CHEMICAL
CONTRACEPTIVES




Julius Schmid, Inc.



The word "RAMSES" is a registered trademark of Julius Schmid, Inc.





"Allergy is perhaps the commonest cause of a stuffy nose...", according to Dill. Shambaugh states that "... 70% of all chronic suppurative sinusitis is on an allergic basis." He emphasizes the importance of controlling the allergic factor in nasal congestion.

Logical therapy for nasal congestion is with this synergistic combination of Antistine, to block the congestive action of histamine, and Privine, to shrink the nasal mucosa. Dosage: 2 to 3 drops in each nostril 3 or 4 times daily.

 Postgrad, Med. 4:413,1948. • 2. M. Rec & Ann. 42:673, 1948.

Antistine®-Privine,® aqueous solution of Antistine (antazoline) hydrochloride 0.5%, and Privine (naphazoline) hydrochloride 0.025%, in bottles of 1 fl. oz. with dropper.

Ciba PHARMACEUTICAL PRODUCTS, INC., SUMMIT, N. J. 2/1632M

#### Antistine-Privine

Antihistaminic plus Vasoconstrictor